

## **Product** Data Sheet

# Decyl maltose neopentyl glycol

 Cat. No.:
 HY-138887

 CAS No.:
 1257852-99-5 

 Molecular Formula:
  $C_{43}H_{80}O_{22}$ 

Molecular Weight: 949.08

Target: Biochemical Assay Reagents

Pathway: Others

**Storage:** 4°C, sealed storage, away from moisture

\* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)

### **SOLVENT & SOLUBILITY**

In Vitro

DMSO: 100 mg/mL (105.37 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	1.0537 mL	5.2683 mL	10.5365 mL
	5 mM	0.2107 mL	1.0537 mL	2.1073 mL
	10 mM	0.1054 mL	0.5268 mL	1.0537 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility:  $\geq$  2.5 mg/mL (2.63 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE- $\beta$ -CD in saline) Solubility: 2.5 mg/mL (2.63 mM); Suspended solution; Need ultrasonic
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (2.63 mM); Clear solution

#### **BIOLOGICAL ACTIVITY**

Description	Decyl maltose neopentyl glycol (DMNG) is the neopentyl glycol detergent that does not disrupt the AlkB oligomeric state. AlkB is a nonheme di-iron alkane hydroxylase <sup>[1]</sup> .
In Vitro	Decyl maltose neopentyl glycol (DMNG) preserves the P. putida AlkB (PpAlkB-SII) hydroxylase activity after purification <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### **REFERENCES**

1]. Hernan Alonso, et al. Chara Pseudomonas putida GPo1. A			e component AlkB of the medium-chain alk	ane hydroxylase system from
	Caution: Product has r	not been fully validated for mo	edical applications. For research use o	nly.
	Tel: 609-228-6898	Fax: 609-228-5909	E-mail: tech@MedChemExpress.c	om
	Address:	1 Deer Park Dr, Suite F, Monmo	outh Junction, NJ 08852, USA	

Page 2 of 2 www.MedChemExpress.com