## L-Glutamic acid-d<sub>3</sub>

Cat. No.:	HY-14608S8	
CAS No.:	203805-84-9	
Molecular Formula:	C <sub>5</sub> H <sub>6</sub> D <sub>3</sub> NO <sub>4</sub>	
Molecular Weight:	150.15	
Target:	Apoptosis; iGluR; Ferroptosis; Endogenous Metabolite	
Pathway:	Apoptosis; Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease	
Storage:	-20°C, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)	

BIOLOGICAL ACTIVITY		
DIOLOGICALACTI		
Description	L-Glutamic acid-d <sub>3</sub> is the deuterium labeled L-Glutamic acid. L-Glutamic acid acts as an excitatory transmitter and an agonist at all subtypes of glutamate receptors (metabotropic, kainate, NMDA, and AMPA). L-Glutamic acid shows a direct activating effect on the release of DA from dopaminergic terminals.	
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

E-mail: tech@MedChemExpress.com

## Product Data Sheet

