Inhibitors

MCE MedChemExpress

Repaglinide M2-d₅

Cat. No.: HY-154749S

 $\label{eq:molecular} \textbf{Molecular Formula:} \qquad C_{27} H_{31} D_5 N_2 O_6$


Molecular Weight: 489.62

Target: Isotope-Labeled Compounds

Pathway: Others

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.



BIOLOGICAL ACTIVITY

Description	Repaglinide M2-d ₅ is deuterated labeled Repaglinide M2.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA