


## H-1152

Cat. No.:	HY-15720
CAS No.:	451462-58-1
Molecular Formula:	C <sub>16</sub> H <sub>21</sub> N <sub>3</sub> O <sub>2</sub> S
Molecular Weight:	319.42
Target:	ROCK
Pathway:	Cell Cycle/DNA Damage; Cytoskeleton; Stem Cell/Wnt; TGF-beta/Smad
Storage:	<div> <div>Powder</div> <div>-20°C    3 years</div> <div>4°C    2 years</div> </div> <div> <div>In solvent</div> <div>-80°C    6 months</div> <div>-20°C    1 month</div> </div>



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (313.07 mM; Need ultrasonic)					
	Preparing Stock Solutions	<div><div>Solvent</div><div>Concentration</div></div>	Mass	1 mg	5 mg	10 mg
		1 mM		3.1307 mL	15.6534 mL	31.3067 mL
		5 mM		0.6261 mL	3.1307 mL	6.2613 mL
		10 mM		0.3131 mL	1.5653 mL	3.1307 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (7.83 mM); Clear solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.5 mg/mL (7.83 mM); Clear solution; Need ultrasonic					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: 2.5 mg/mL (7.83 mM); Clear solution; Need ultrasonic					

### BIOLOGICAL ACTIVITY

Description	H-1152 is a membrane-permeable and selective ROCK inhibitor, with a K <sub>i</sub> value of 1.6 nM, and an IC <sub>50</sub> value of 12 nM for ROCK2.			
IC <sub>50</sub> & Target	ROCKII 12 nM (IC <sub>50</sub> )	CaMKII 0.18 μM (IC <sub>50</sub> )	PKG 0.36 μM (IC <sub>50</sub> )	AuroraA 0.745 μM (IC <sub>50</sub> )
	PKA 3.03 μM (IC <sub>50</sub> )	Src 3.06 μM (IC <sub>50</sub> )	PKC 5.68 μM (IC <sub>50</sub> )	Abl 7.77 μM (IC <sub>50</sub> )

	MKK4 16.9 $\mu$ M (IC <sub>50</sub> )	MLCK 28.3 $\mu$ M (IC <sub>50</sub> )	EGFR 50 $\mu$ M (IC <sub>50</sub> )	GSK3 $\alpha$ 60.7 $\mu$ M (IC <sub>50</sub> )
	AMPK 100 $\mu$ M (IC <sub>50</sub> )	P38 $\alpha$ 100 $\mu$ M (IC <sub>50</sub> )		
<b>In Vitro</b>	<p>H-1152 is an inhibitor of Rho-kinase, with an IC<sub>50</sub> of 12 nM for ROCK2. H-1152 (H-1152P) also shows less inhibitory activities against CaMKII, PKG, AuroraA, PKA, Src, PKC, MLCK, Abl, EGFR, MKK4, GSK3<math>\alpha</math>, AMPK, and P38<math>\alpha</math>, with IC<sub>50</sub>s of 0.180, 0.360, 0.745, 3.03, 3.06, 5.68, 28.3, 7.77, 50.0, 16.9, 60.7, 100, and 100 <math>\mu</math>M, respectively<sup>[1]</sup>. H-1152 potently inhibits Rho kinase, with a K<sub>i</sub> of 1.6 nM, and slightly suppresses PKA, PKC and MLCK, with K<sub>i</sub>s of 0.63, 9.27, and 10.1 <math>\mu</math>M, respectively. H-1152 (0.1-10 <math>\mu</math>M) highly inhibits MARCKS phosphorylation, with an IC<sub>50</sub> value of 2.5 <math>\mu</math>M in LPA-treated cells, but shows no such obvious effects in PDBu-treated cells<sup>[2]</sup>. H-1152 (0.5-10 <math>\mu</math>M) causes no decreased neuronal survival. H-1152 (1, 5 or 10 <math>\mu</math>M) also exerts no alterations in the ratios of different neuronal morphologies. Furthermore, H-1152 (10 <math>\mu</math>M) increases neurite length in both BMP4 and LIF cultures<sup>[3]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>			

## PROTOCOL

### Kinase Assay <sup>[2]</sup>

Inhibitors (including H-1152) are added at the indicated concentrations to 50  $\mu$ L of the assay mixture 50 mM Tris-HCl (pH 7.5), 5 mM MgCl<sub>2</sub>, 1 mM EDTA, 1 mM EGTA, 1 mM dithiothreitol, 40  $\mu$ M S6-peptide, various concentrations of [ $\gamma$ -<sup>32</sup>P]ATP and purified Rho-kinase. The reactions are started by the addition of [ $\gamma$ -<sup>32</sup>P]ATP and carried out at 30°C for 5 min. The Michaelis-Menten equation is used to calculate the Michaelis constant (K<sub>m</sub>) and maximal velocity (V<sub>max</sub>) of Rho-kinase. Data are further analyzed with secondary plot to calculate the inhibitory constant (K<sub>i</sub>)<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Cell Assay <sup>[3]</sup>

Briefly, cells are routinely plated on poly-d-lysine/laminin coated 96 well plates or in 16 well glass culture slides. Control medium contained Dulbecco's modified Eagles medium/Hams F12(1:1) (DMEM/F12), 2 mM L-glutamine, N2 mix (1:100 dilution), 0.63 mL of 45% glucose for each 100 mL of DMEM/F12, neurotrophin 3 (NT3; final concentration, 8 ng/mL), BDNF (final concentration 8 ng/mL), and 10% fetal bovine serum heat inactivated before use. LIF cultures contain control medium+LIF (50 ng/mL). BMP4 cultures contain control medium+bone morphogenetic protein 4 (BMP4; 25 ng/mL). Total volume of culture is 110  $\mu$ L. ROCK inhibitor H-1152 is diluted in water and added in an additional 10  $\mu$ L to cultures 24 h after plating. Water is added to controls. Eighteen hours after the addition of inhibitor, cultures are fixed in 4% paraformaldehyde (1 h at room temperature for peroxidase-linked labeling and 20 min at room temperature for fluorescence labeling). For ArrayScan/Cellomics automated analysis: Cells are plated in a total volume of 50  $\mu$ L on 384 well plastic plates previously coated with poly-d-lysine/laminin, and cultured in the same medium<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Research Square Preprint. 2021 Jun.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

- [1]. Tamura M, et al. Development of specific Rho-kinase inhibitors and their clinical application. Biochim Biophys Acta. 2005 Dec 30;1754(1-2):245-52. Epub 2005 Sep 12.
- [2]. Ikenoya M, et al. Inhibition of rho-kinase-induced myristoylated alanine-rich C kinase substrate (MARCKS) phosphorylation in human neuronal cells by H-1152, a novel and specific Rho-kinase inhibitor. J Neurochem. 2002 Apr;81(1):9-16.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA