


## Polidocanol

Cat. No.:	HY-B2106
CAS No.:	9002-92-0
Molecular Formula:	$(C_2H_4O)_n C_{12}H_{26}O$
Target:	Others
Pathway:	Others
Storage:	<div>Pure form</div> <div>-20°C 3 years</div> <div>4°C 2 years</div> <div>In solvent</div> <div>-80°C 6 months</div> <div>-20°C 1 month</div>



### SOLVENT & SOLUBILITY

In Vitro	H <sub>2</sub> O : 50 mg/mL (Need ultrasonic) DMSO : 50 mg/mL (Need ultrasonic)
In Vivo	1. Add each solvent one by one: PBS Solubility: 120 mg/mL (Infinity mM); Clear solution; Need ultrasonic

### BIOLOGICAL ACTIVITY

Description	Polidocanol is a sclerosing agent used successfully to treat extremity and esophageal varices and telangiectasias.
In Vivo	Polidocanol (0.5%, 1%, and 2%) results in significantly more adhesions than tetracycline, whereas the effect of diluted polidocanol is similar to that of tetracycline in rats. The higher concentrations of polidocanol, of 0.5%, 1%, and 2% are more effective than either the diluted polidocanol or tetracycline against pleurodesis <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Cetin B, et al. Polidocanol at different concentrations for pleurodesis in rats. Surg Today. 2005;35(12):1066-9.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA