


## Calycosin-7-O-β-D-glucoside

Cat. No.:	HY-N0520
CAS No.:	20633-67-4
Molecular Formula:	C <sub>22</sub> H <sub>22</sub> O <sub>10</sub>
Molecular Weight:	446.4
Target:	Reactive Oxygen Species
Pathway:	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB
Storage:	Powder    -20°C    3 years 4°C    2 years In solvent   -80°C    2 years -20°C    1 year



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 100 mg/mL (224.01 mM)

\* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.2401 mL	11.2007 mL	22.4014 mL
	5 mM		0.4480 mL	2.2401 mL	4.4803 mL
	10 mM		0.2240 mL	1.1201 mL	2.2401 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (5.60 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (5.60 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 2.5 mg/mL (5.60 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Calycosin-7-O-β-D-glucoside is an isoflavone isolated from Astragali Radix. Calycosin-7-O-β-D-glucoside has variety of biological activities, such as neuroprotective, cardioprotection, anti-inflammation, and antioxidative stress effects<sup>[1][2]</sup>.

#### In Vitro

Calycosin-7-O-β-D-glucoside (2 μM; 6 hours) remarkably inhibits the expression and activities of MMPs, and secures the expression of cav-1 and tight junction proteins in the microvessels isolated from ischemic rat cortex<sup>[1]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## In Vivo

Calycosin-7-O- $\beta$ -D-glucoside (intraperitoneal injection; 26.8 mg/kg; 14 days) significantly reduces infarct volume, histological damage and BBB permeability in the in vivo MCAO ischemia-reperfusion rat model<sup>[1]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Middle cerebral artery occlusion (MCAO) male adult Sprague-Dawley rats <sup>[1]</sup>
Dosage:	26.8 mg/kg
Administration:	Intraperitoneal injection; 26.8 mg/kg; 14 days
Result:	Exhibited neuroprotective effects in rats.

## CUSTOMER VALIDATION

- Acta Pharm Sin B. 2021 Jan;11(1):143-155.
- Chem Biol Interact. 2023 Feb 20;110411.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

- [1]. Shuping Fu, et al. Calycosin-7-O- $\beta$ -D-glucoside regulates nitric oxide /caveolin-1/matrix metalloproteinases pathway and protects blood-brain barrier integrity in experimental cerebral ischemia-reperfusion injury. J Ethnopharmacol. 2014 Aug 8;155(1):692-701.
- [2]. Xiangli Yan, et al. Calycosin-7- O-  $\beta$ - D-glucoside Attenuates OGD/R-Induced Damage by Preventing Oxidative Stress and Neuronal Apoptosis via the SIRT1/FOXO1/PGC-1  $\alpha$  Pathway in HT22 Cells. Neural Plast. 2019 Dec 1;2019:8798069.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA