Furaneol-¹³C₂

\cap
$\sqrt{\frac{13}{13}}$
O'SC
¹³ CH ₃

Product Data Sheet

BIOLOGICAL ACTIV	
Description	Furaneol- $^{13}C_2$ is ^{13}C labeled Nonanal (HY-N8016). Nonanal is a saturated fatty aldehyde with antidiarrhoeal activity ^[1] .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Sasaki K, et al. Molecular cloning and characterization of UDP-glucose: furaneol glucosyltransferase gene from grapevine cultivar Muscat Bailey A (Vitis labrusca × V. vinifera. J Exp Bot. 2015 Oct;66(20):6167-74.

[2]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

MedChemExpress

Tel: 609-228-6898