

Product Data Sheet

Bombesin

 Cat. No.:
 HY-P0195

 CAS No.:
 31362-50-2

 Molecular Formula:
 $C_{71}H_{110}N_{24}O_{18}S$

Molecular Weight: 1620

Sequence: {Glp}-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leu-Met-NH2

Sequence Shortening: {Glp}-QRLGNQWAVGHLM-NH2

Target: Bombesin Receptor

Pathway: GPCR/G Protein

Storage: Sealed storage, away from moisture and light

Powder -80°C 2 years -20°C 1 year

* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture

and light)

SOLVENT & SOLUBILITY

In Vitro

H₂O: 50 mg/mL (30.86 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	0.6173 mL	3.0864 mL	6.1728 mL
	5 mM	0.1235 mL	0.6173 mL	1.2346 mL
	10 mM	0.0617 mL	0.3086 mL	0.6173 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Bombesin, a tetradecapeptide, plays an important role in the release of gastrin and the activation of G-protein receptors [1].

In Vitro

Bombesin is a tetradecapeptide with a COOH terminus ending in Gly-His-Leu-Met-NH2 and subsequently is shown to closely resemble two mammalian bombesin -related peptides, gastrin-releasing peptide (GRP) and neuromedin B (NMB)^[1]. Bombesin is found to have stimulatory effects upon gastric and pancreatic secretions, release of gastrointestinal hormones, gallbladder contraction and bronchoconstriction. It is present in amphibian gastric endocrine cells, avian proventriculus endocrine cells and avian brain. In mammals it is present mainly in nerve cells and fibers. The only mammalian endocrine cell shown to date to have bombesin is the P-cell in fetal lung. Bombesin is also found in mammalian brain, with its highest concentration in the hypothalamus^[2]. Bombesin is shown to be a potent mitogen for Swiss 3T3 cells. In the presence of a low concentration (3.5%) of serum, bombesin stimulates 3T3 cell proliferation. In serum-free medium, bombesin induces DNA synthesis in the absence of any other added growth factor (IC₅₀=1 nM)^[3].

 $\label{eq:mce} \mbox{MCE has not independently confirmed the accuracy of these methods. They are for reference only.}$

CUSTOMER VALIDATION

• Brain Res. 2022 May 23;1789:147950.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Gonzalez N, et al. Bombesin-related peptides and their receptors: recent advances in their role in physiology and disease states. Curr Opin Endocrinol Diabetes Obes. 2008 Feb;15(1):58-64.
- [2]. Chejfec G, et al. Bombesin in human neuroendocrine (NE) neoplasms. Peptides. 1985;6 Suppl 3:107-12.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

 $\hbox{E-mail: } tech @ Med Chem Express.com$

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA