


Histone H3K9me3 (1-15) (TFA)

Cat. No.:	HY-P10111A
Molecular Formula:	$C_{66}H_{124}N_{25}O_{21} \cdot xC_2HF_3O_2$
Sequence:	Ala-Arg-Thr-Lys-Gln-Thr-Ala-Arg-{Lys(Me3)}-Ser-Thr-Gly-Gly-Lys-Ala
Sequence Shortening:	ARTKQTAR-{Lys(Me3)}-STGGKA
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description

Histone H3K9me3 (1-15) (H3(1-15)K9me3) TFA is used as substrate. Histone H3K9me3 is a histone posttranslational modification (PTM) that has emerged as hallmark of pericentromeric heterochromatin^{[1][2]}.

REFERENCES

- [1]. Kerstin Mosch, et al. HP1 recruits activity-dependent neuroprotective protein to H3K9me3 marked pericentromeric heterochromatin for silencing of major satellite repeats. PLoS One. 2011 Jan 18;6(1):e15894.
- [2]. Esther C Y Woon, et al. Linking of 2-oxoglutarate and substrate binding sites enables potent and highly selective inhibition of JmjC histone demethylases. Angew Chem Int Ed Engl. 2012 Feb 13;51(7):1631-4.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA