

## CD200 Protein, Cynomolgus (HEK293, His)

Cat. No.:	HY-P700670
Synonyms:	CD200; MOX1; MOX2; My033; MRC; OX-2; OX2
Species:	Cynomolgus
Source:	HEK293
Accession:	A0A2K5TQS2 (Q56-G257)
Gene ID:	/
Molecular Weight:	45-55 kDa

### PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 $\mu$ m filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/ $\mu$ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	The CD200 protein plays a pivotal role in immune modulation, functioning as a costimulatory molecule that promotes T-cell proliferation. Additionally, CD200 is implicated in the potential regulation of myeloid cell activity across various tissues, as suggested by similarity-based inferences. The interaction between CD200 and its receptor, CD200R1, is facilitated through their respective N-terminal Ig-like domains. This interaction likely contributes to the regulatory processes involved in immune responses and myeloid cell function. The engagement of CD200 with CD200R1 underscores its significance in mediating immune cell communication and homeostasis, emphasizing its potential as a key player in immunomodulation.
------------	--

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA