

Legumain Protein, Mouse (HEK293, C-His)

Cat. No.:	HY-P70216A
Synonyms:	rMuLegumain/Asparaginyl Endopeptidase, His; Legumain; Lgmn; Asparaginyl endopeptidase; Protease cysteine 1; Prsc1
Species:	Mouse
Source:	HEK293
Accession:	O89017 (V18-Y435)
Gene ID:	19141
Molecular Weight:	Approximately 55-60 kDa

PROPERTIES

AA Sequence	<pre> V P V G V D D P E D G G K H W V V I V A G S N G W Y N Y R H Q A D A C H A Y Q I I H R N G I P D E Q I I V M M Y D D I A N S E E N P T P G V V I N R P N G T D V Y K G V L K D Y T G E D V T P E N F L A V L R G D A E A V K G K G S G K V L K S G P R D H V F I Y F T D H G A T G I L V F P N D D L H V K D L N K T I R Y M Y E H K M Y Q K M V F Y I E A C E S G S M M N H L P D D I N V Y A T T A A N P K E S S Y A C Y Y D E E R G T Y L G D W Y S V N W M E D S D V E D L T K E T L H K Q Y H L V K S H T N T S H V M Q Y G N K S I S T M K V M Q F Q G M K H R A S S P I S L P P V T H L D L T P S P D V P L T I L K R K L L R T N D V K E S Q N L I G Q I Q Q F L D A R H V I E K S V H K I V S L L A G F G E T A E R H L S E R T M L T A H D C Y Q E A V T H F R T H C F N W H S V T Y E H A L R Y L Y V L A N L C E A P Y P I D R I E M A M D K V C L S H Y </pre>
Biological Activity	<ol style="list-style-type: none"> 1. Measured in a cell proliferation assay using RAW264.7 cells. The ED₅₀ this effect is 0.07438ng/mL, corresponding to a specific activity is 1.344×10⁷ units/mg. 2. Measured by its ability to cleave the fluorogenic peptide substrate, N-carbobenzyloxy-Ala-Ala-Asn-7-amido-4-methylcoumarin (Z-AAN-AMC). The specific activity is >130 pmol/min/μg, as measured under the described conditions.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μm filtered solution of 50 mM Tris-HCL/Tris, 300 mM NaCl, pH 7.4.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Legumain protein exhibits a strict specificity for the hydrolysis of asparaginyl bonds. Additionally, it demonstrates the ability to cleave aspartyl bonds slowly, particularly in acidic conditions, further expanding its enzymatic versatility. Functionally, Legumain is integral to the processing of proteins for MHC class II antigen presentation within the lysosomal/endosomal system. It also plays a crucial role in MHC class I antigen presentation in cross-presenting dendritic cells by facilitating the cleavage and maturation of Perforin-2 (MPEG1), thereby promoting antigen translocation in the cytosol, as indicated by recent research findings. Moreover, Legumain is essential for normal lysosomal protein degradation in renal proximal tubules and is required for the degradation of internalized EGFR, highlighting its importance in cellular processes and the regulation of cell proliferation.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA