

## Product Data Sheet

## CDK2-CCNO Protein, Human (sf9, GST, Flag, His)

Cat. No.:	HY-P702679
Synonyms:	CDK2; cyclin-dependent kinase 2; p33 protein kinase; cell devision kinase 2; cdc2-related protein kinase; cell division protein kinase 2; p33(CDK2)
Species:	Human
Source:	Sf9 insect cells
Accession:	P24941-1 (M1-L298)&P22674-1 (V2-S349)
Gene ID:	1017&10309
Molecular Weight:	60.5 kDa&40.7 kDa

ProtectionBiological ActivityThe activity was measured by off-chip mobility shift assay(MSA). The enzyme was incubated with fluorecence-labeled substrate and Mg(or Mn)/ATP. The phosphorylated and unphosphorylated substrates were separated and detected by MSA device. The Km of CDK2-CCN0 for its substrate is 95.74 µM.AppearanceSolution.FormulationSupplied as a 0.2 µm filtered solution of 50 mM HEPES (pH 7.5), 200 mM NaCl, 5% glycerol, 1 mM DTT.Endotoxin Level<1 EU/µg, determined by LAL method.ReconsititutionPlease use rapid thawing with running water to thaw the protein.Storage & StabilityStored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.ShippingShipping with dry ice.	PROPERTIES	
Substrate and Mg(or Mn)/ATP. The phosphorylated and unphosphorylated substrates were separated and detected by MSA device. The Km of CDK2-CCNO for its substrate is 95.74 µM.AppearanceSolution.FormulationSupplied as a 0.2 µm filtered solution of 50 mM HEPES (pH 7.5), 200 mM NaCl, 5% glycerol, 1 mM DTT.Endotoxin Level<1 EU/µg, determined by LAL method.	PROPERTIES	
Formulation Supplied as a 0.2 μm filtered solution of 50 mM HEPES (pH 7.5), 200 mM NaCl, 5% glycerol, 1 mM DTT.   Endotoxin Level <1 EU/μg, determined by LAL method.	Biological Activity	substrate and Mg(or Mn)/ATP. The phosphorylated and unphosphorylated substrates were separated and detected by MSA
Endotoxin Level <1 EU/µg, determined by LAL method.	Appearance	Solution.
Reconsititution Please use rapid thawing with running water to thaw the protein.   Storage & Stability Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.	Formulation	Supplied as a 0.2 $\mu m$ filtered solution of 50 mM HEPES (pH 7.5), 200 mM NaCl, 5% glycerol, 1 mM DTT.
Storage & Stability Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.	Endotoxin Level	<1 EU/µg, determined by LAL method.
extended storage. Avoid repeated freeze-thaw cycles.	Reconsititution	Please use rapid thawing with running water to thaw the protein.
Shipping Shipping with dry ice.	Storage & Stability	
	Shipping	Shipping with dry ice.

## Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609	228-5909
----------------------------	----------

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA