

Uteroglobin/SCGB1A1 Protein, Human (HEK293, His)

Cat. No.:	HY-P71107
Synonyms:	Uteroglobin; Clara cell phospholipid-binding protein; CCPBP; Clara cells 10 kDa secretory protein; CC10; Secretoglobin family 1A member 1; Urinary protein 1; UP-1; UP1; Urine protein 1; SCGB1A1; CCSP; UGB
Species:	Human
Source:	HEK293
Accession:	P11684 (E22-N91)
Gene ID:	7356
Molecular Weight:	Approximately 9.0 kDa

PROPERTIES

AA Sequence	E I C P S F Q R V I E T L L M D T P S S Y E A A M E L F S P D Q D M R E A G A Q L K K L V D T L P Q K P R E S I I K L M E K I A Q S S L C N
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	The Uteroglobin/SCGB1A1 protein exhibits versatile binding capabilities, including the ability to bind phosphatidylcholine, phosphatidylinositol, polychlorinated biphenyls (PCB), and weakly binds progesterone. Additionally, it functions as a potent inhibitor of phospholipase A2. Structurally, this protein forms an antiparallel homodimer that is disulfide-linked. While there is some controversy surrounding its interaction with LMBR1L, further investigation is required to clarify this aspect.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA