

HA/Hemagglutinin Protein, H7N9 (EPI439507, sf9, His)

Cat. No.:	HY-P73963
Synonyms:	HA; Hemagglutinin; HA/Hemagglutinin Protein, H7N9 (A/Anhui/1/2013, sf9, His)
Species:	Virus
Source:	Sf9 insect cells
Accession:	EPI439507 (D19-V524)
Gene ID:	/
Molecular Weight:	Approximately 58 kDa

PROPERTIES

AA Sequence	<pre> DKICLGHHAV SNGTKVNTLT ERGVEVVNAT ETVERTNIPR ICSKGKRTVD LGQCGLLGTI TGPPQCDQFL EFSADLIIER REGSDVCYPG KFNVEEALRQ ILRESGGIDK EAMGFTYSGI RTNGATSACR RSGSSFYAEM KWLLSNTDNA AFPQMTKSYK NTRKSPALIV WGIHHSVSTA EQTKLYGSGN KLVTVGSSNY QQSFVPSPGA RPQVNGLSGR IDFHWMMLNP NDTVTF SFNG AFIAPDRASF LRGKSMGIQS GVQVDANCEG DCYHSGGTII SNLPPFNIDS RAVGKCPRYV KQRSLLLATG MKNVPEIPKG RGLFGAIAGF IENGWEGLID GWYGFRRHQA QGEGTAADYK STQSAIDQIT GKLNRLIEKT NQQFELIDNE FNEVEKQIGN VINWTRDSIT EVWSYNAELL VAMENQHTID LADSEMDKLY ERVKRQLREN A EEDGTGCFE IFHKCDDDCM ASIRNNTYDH SKYREEAMQN RIQIDPVKLS SGYKDV </pre>
Appearance	Solution.
Formulation	Supplied as a 0.22 µm filtered solution of 20 mM Tris, 500 mM NaCl, 10% glycerol, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

DESCRIPTION

Background

The Hemagglutinin (HA) protein plays a crucial role in the attachment of virus particles to host cells by binding to sialic acid-containing receptors on the cell surface. This interaction not only induces virion internalization through clathrin-dependent endocytosis but also facilitates an alternative clathrin- and caveolin-independent pathway for about one-third of the virus particles. HA is a Class I viral fusion protein responsible for penetrating the cell cytoplasm by mediating the fusion of the virus particle's membrane with the endosomal membrane. The low pH environment in endosomes triggers an irreversible conformational change in HA2, leading to the release of the fusion hydrophobic peptide. The cooperative action of several trimers is necessary to form a competent fusion pore, highlighting the intricate role of HA in host range restriction and virulence determination.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA