

ACVR2A/Activin RIIA Protein, Mouse (HEK293, His)

Cat. No.:	HY-P75563
Synonyms:	ACVR-2A; Activin receptor type 2A; ACTR-IIA; ACVR2
Species:	Mouse
Source:	HEK293
Accession:	P27038 (A20-P134)
Gene ID:	11480
Molecular Weight:	Approximately 35 kDa

PROPERTIES

AA Sequence	<p>A I L G R S E T Q E C L F F N A N W E R D R T N Q T G V E P C Y G D K D K R R H</p> <p>C F A T W K N I S G S I E I V K Q G C W L D D I N C Y D R T D C I E K K D S P E</p> <p>V Y F C C C E G N M C N E K F S Y F P E M E V T Q P T S N P V T P K P</p>
Biological Activity	Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED ₅₀ for this effect is typically 0.4-3 µg/mL in the presence of 10 ng/mL Recombinant Human Activin A.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>ACVR2A is a type II member of the TGF-β family of receptor Serine/Threonine kinases. ACVR2A is a receptor for Activin A, Activin B and Inhibin A^{[1][2]}.</p> <p>The sequence of amino acids in ACVR2A proteins from different species is very stable, which leads to the conclusion that in the process of evolution, ACVR2A has been only slightly altered, and that both in humans and in animals, its function is similar.</p> <p>Signaling by activins and BMPs is highly promiscuous, since apart from signaling through ALK4/7, the activin type II</p>
------------	--

receptors (ACVR2A and 2B) can interact also with several type I BMP receptors (ALK1/2/3/6), which can also form complexes with the type II BMP receptor, BMPRII. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. ACVR2A can form complexes with different type I receptors that signal either to Smad2/3 (ALK4) or to Smad1/5/8 (ALK2, ALK3, ALK6). The different type I receptors compete for binding to ACVR2A and that this competition provides a mechanism that balances signaling between Activin A-mediated, ALK4-dependent Smad2/3 signaling, and BMP-mediated ALK2 or ALK3-dependent signaling to Smad1/5/8. In myeloma cells, BMP-6- and BMP-9-induced activation of SMAD1/5/8 through ACVR2A/ACVR2B/ALK2 is inhibited by activin A treatment^{[1][3]}.

REFERENCES

- [1]. Oddrun Elise Olsen, et al. Activin A inhibits BMP-signaling by binding ACVR2A and ACVR2B. *Cell Commun Signal*. 2015 Jun 6;13:27.
- [2]. V K Chin, et al. Inhibition of Activin A suppressed tumor necrosis factor- α secretion and improved histopathological conditions in malarial mice. *Trop Biomed*. 2021 Mar 1;38(1):187-204.
- [3]. Szabina Szófia Szilágyi, et al. Competition between type I activin and BMP receptors for binding to ACVR2A regulates signaling to distinct Smad pathways. *BMC Biol*. 2022 Feb 18;20(1):50.
- [4]. Heather L Hayes, et al. A Pdx-1-Regulated Soluble Factor Activates Rat and Human Islet Cell Proliferation. *Mol Cell Biol*. 2016 Nov 14;36(23):2918-2930.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA