

ACE2 Protein, Human (Biotinylated, HEK293, Fc-Avi)

Cat. No.:	HY-P78055
Synonyms:	ACE-2; ACEH; ACE2
Species:	Human
Source:	HEK293
Accession:	Q9BYF1 (Q18-S740)
Gene ID:	59272
Molecular Weight:	115-125 kDa

PROPERTIES

Biological Activity	Immobilized SARS-CoV-2 Spike S at 2 µg/mL (100µL/Well) on the plate. Dose response curve for Biotinylated Human ACE2-hFc with the EC ₅₀ of 0.3-0.7 ug/ml determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

ACE2, an indispensable counter-regulatory carboxypeptidase within the renin-angiotensin hormone system, plays a pivotal role in maintaining cardiovascular homeostasis by intricately regulating blood volume and systemic vascular resistance. Through its enzymatic activity, ACE2 converts angiotensin I to angiotensin 1-9 and angiotensin II to angiotensin 1-7, exerting anti-hypertrophic effects in cardiomyocytes and acting as a vasodilator with anti-proliferative properties. Beyond its central role in the renin-angiotensin system, ACE2 exhibits broad enzymatic activity, cleaving various vasoactive peptides such as neurotensin, kinetensin, and des-Arg bradykinin. Moreover, ACE2 is proficient in cleaving other biological peptides, including apelin, casomorphins, and dynorphin A. Notably, ACE2's C-terminus, homologous to collectrin, orchestrates the trafficking of the neutral amino acid transporter SL6A19 to the gut epithelial cell membrane, thereby regulating its surface expression and catalytic activity. Importantly, ACE2 also serves as a receptor for human coronaviruses SARS-CoV, SARS-CoV-2, and HCoV-NL63, implicating it in microbial infection pathways.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA