



## **Product** Data Sheet

## Semaphorin-4D/SEMA4D Protein, Human (HEK293, His-Avi)

Cat. No.: HY-P78512

Semaphorin-4D; CD100; SEMA4D; C9orf164; FLJ33485; FLJ34282; FLJ39737; FLJ46484; M-sema-Synonyms:

G; MGC169138; MGC169141; SEMAJ; coll-4

Species: Human **HEK293** Source:

Accession: Q92854 (M22-R734)

Gene ID: 10507

Molecular Weight: 115-140 kDa

|  |  |  | B |
|--|--|--|---|
|  |  |  |   |
|  |  |  |   |
|  |  |  |   |

| Biological Activity | Immobilized Human Semaphorin 4D, His Tag at $1\mu g/ml$ ( $100\mu l/well$ ) on the plate. Dose response curve for Anti-Semaphorin 4D Antibody , hFc Tag with the EC <sub>50</sub> of 9.3ng/ml determined by ELISA. |  |  |
|---------------------|--|--|--|
| Appearance          | Lyophilized powder.  |  |  |
| Formulation         | Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.   |  |  |
| Endotoxin Level     | <1 EU/μg, determined by LAL method.  |  |  |
| Reconsititution     | It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH <sub>2</sub> O.   |  |  |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.         |  |  |
| Shipping            | Room temperature in continental US; may vary elsewhere.  |  |  |

## **DESCRIPTION**

## **Background**

Semaphorin-4D/SEMA4D Protein, a cell surface receptor for PLXNB1 and PLXNB2, plays a crucial role in cell-cell signaling. It is involved in regulating GABAergic synapse development and promotes the development of inhibitory synapses in a PLXNB1-dependent manner. By activating PLXNB1 and interacting with it, SEMA4D modulates the complexity and arborization of developing neurites in hippocampal neurons, mediating the activation of RHOA. Additionally, SEMA4D promotes the migration of cerebellar granule cells and induces B-cells to aggregate, enhancing their viability in vitro. In the immune system, SEMA4D plays a role in inducing endothelial cell migration through the activation of PTK2B/PYK2, SRC, and the phosphatidylinositol 3-kinase-AKT pathway. It forms homodimers and interacts with PLXNB2, as well as interacts with PLXNB1.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com