

## CEACAM1 Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P78894
Synonyms:	CEACAM1; CD66a; BGP; BGP1; BGPI
Species:	Human
Source:	HEK293
Accession:	P13688 (Q35-G428)
Gene ID:	634
Molecular Weight:	66-115 kDa

### PROPERTIES

Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized Biotinylated Human CEACAM1 at 1 µg/mL on streptavidin coated plats can bind Anti-CEACAM1 antibody. The EC <sub>50</sub> is 0.3405-0.6093 ng/mL.
Appearance	Lyophilized powder
Formulation	Lyophilized a 0.22 µm filtered solution of PBS, 6% Trehalose, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	The SF3R gene is involved in various biological processes and functions. It regulates insulin action by promoting the clearance of insulin and regulating lipogenesis in the liver. Upon insulin stimulation, SF3R undergoes phosphorylation by the insulin receptor (INSR), leading to increased insulin endocytosis and degradation. This results in a reduction of fatty acid synthesis. SF3R also plays a role in down-regulating cell proliferation through its interaction with SHC1, which decreases the coupling of SHC1 to the MAPK3/ERK1-MAPK1/ERK2 and phosphatidylinositol 3-kinase pathways.
------------	---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA