MCE RedChemExpress

Guanine-13C,15N2

 Cat. No.:
 HY-Y1055S1

 CAS No.:
 202407-12-3

 Molecular Formula:
 C₄13CH₅N₃15N₃O

Molecular Weight: 154.11

Target: DNA/RNA Synthesis; Endogenous Metabolite

Pathway: Cell Cycle/DNA Damage; Metabolic Enzyme/Protease

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

$$H_{2}N$$
 $H_{2}N$
 $H_{3}CH$
 $H_{2}N$
 $H_{3}CH$

BIOLOGICAL ACTIVITY

Description	Guanine- 13 C, 15 N ₂ is the 13 C and 15 N labeled Guanine[1]. Guanine (2-Aminohypoxanthine) is one of the fundamental components of nucleic acids (DNA and RNA). Guanine is a purine derivative, consisting of a fused pyrimidine-imidazole ring system with conjugated double bonds.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA