


Cadazolid

Cat. No.:	HY-100436		
CAS No.:	1025097-10-2		
Molecular Formula:	C ₂₉ H ₂₉ F ₂ N ₃ O ₈		
Molecular Weight:	585.55		
Target:	Bacterial; Antibiotic		
Pathway:	Anti-infection		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (170.78 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
			1 mg	5 mg
			10 mg	
Preparing Stock Solutions	1 mM	1.7078 mL	8.5390 mL	17.0780 mL
	5 mM	0.3416 mL	1.7078 mL	3.4156 mL
	10 mM	0.1708 mL	0.8539 mL	1.7078 mL
Please refer to the solubility information to select the appropriate solvent.				
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (4.27 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (4.27 mM); Clear solution			

BIOLOGICAL ACTIVITY

Description	Cadazolid (ACT-179811) is a new oxazolidinone antibiotic with potent activity against <i>Clostridium difficile</i> .
IC ₅₀ & Target	Oxazolidinone
In Vitro	Cadazolid is a new antibiotic in development for the treatment of <i>Clostridium difficile</i> -associated diarrhea ^[1] . Cadazolid is active against all (including linezolid- and moxifloxacin-resistant) <i>Clostridium difficile</i> strains (MIC ₉₀ 0.125, range 0.03-0.25 mg/L). The cadazolid geometric mean MIC is 152-fold, 16-fold, 9-fold and 7-fold lower than those of moxifloxacin, linezolid, metronidazole and vancomycin, respectively. Both cadazolid dosing regimens rapidly reduce <i>Clostridium difficile</i> viable counts and cytotoxin with no evidence of recurrence. Cadazolid levels persists at 50-100-fold supra-MIC for 14 days post-dosing. Cadazolid inhibition of enumerated gut microflora is limited, with the exception of bifidobacteria; <i>Bacteroides</i>

fragilis group and Lactobacillus spp. counts are unaffected. There is no evidence for selection of strains resistant to cadazolid, quinolones or linezolid^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Cadazolid is well tolerated up to 3000 mg given twice daily for 10 days. The most common adverse event is headache, with no observed relationship between dose or treatment duration and adverse events. Plasma concentrations of cadazolid are low. No plasma concentrations >3.3 ng/mL are observed after single doses or >6.9 ng/mL after 10 days of multiple doses. Food increased the mean C_{max} from 0.73 to 1.87 ng/mL and mean AUC_{0-t} from 3.13 to 15.69 ng-h/mL after a single 300 mg dose. The increase in systemic exposure to cadazolid across doses is less than dose-proportional. The mean cumulative faecal recovery is 81.0%–93.5%. Urinary recovery of unchanged compound is less than 0.015%^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

MICs of cadazolid, metronidazole, vancomycin, moxifloxacin and linezolid are determined using agar incorporation for 100 Clostridium difficile isolates, including 30 epidemic strains (ribotypes 027, 106 and 001) with reduced metronidazole susceptibility, 2 linezolid-resistant isolates and 2 moxifloxacin-resistant isolates. The efficacy of two cadazolid dosing regimens (250 versus 750 mg/L twice daily for 7 days) to treat simulated CDI is evaluated. Microflora populations, Clostridium difficile total viable counts and spores, cytotoxin titres, possible emergence of cadazolid, linezolid or quinolone resistance, and antimicrobial concentrations are monitored throughout^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Baldoni D, et al. Cadazolid, a novel antibiotic with potent activity against Clostridium difficile: safety, tolerability and pharmacokinetics in healthy subjects following single and multiple oral doses. J Antimicrob Chemother. 2014 Mar;69(3):706-14.

[2]. Chilton CH, et al. In vitro activity of cadazolid against clinically relevant Clostridium difficile isolates and in an in vitro gut model of C. difficile infection. J Antimicrob Chemother. 2014 Mar;69(3):697-705.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA