


NOD-IN-1

Cat. No.:	HY-100691		
CAS No.:	132819-92-2		
Molecular Formula:	C ₁₈ H ₁₇ NO ₄ S		
Molecular Weight:	343.4		
Target:	NOD-like Receptor (NLR)		
Pathway:	Immunology/Inflammation		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (291.21 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	2.9121 mL	14.5603 mL	29.1206 mL
		5 mM	0.5824 mL	2.9121 mL	5.8241 mL
10 mM		0.2912 mL	1.4560 mL	2.9121 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.28 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.28 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.28 mM); Clear solution 				

BIOLOGICAL ACTIVITY

Description	NOD-IN-1 is a potent mixed inhibitor of nucleotide-binding oligomerization domain (NOD)-like receptors, NOD1 and NOD2, with IC ₅₀ of 5.74 μM and 6.45 μM, respectively.
IC₅₀ & Target	IC ₅₀ : 5.74 μM (NOD1), 6.45 μM (NOD2) ^[1]
In Vitro	NOD-IN-1 (compound 4) is potent mixed inhibitor of NOD1 and NOD2, displaying a balanced inhibitory activity on both targets in the low micromolar range. NOD-IN-1 (IC ₅₀ (NOD1)=5.74 μM; IC ₅₀ (NOD2)=6.45 μM) is identified as the best of the

series, possessing NOD1- and NOD2-inhibitory activities in the lower micromolar range. These results show that NOD-IN-1 is 7-fold less potent than Noditinib-1 in terms of NOD1 inhibition and completely devoid of selective activity for NOD1 or NOD2 as opposed to Noditinib-1. NOD-IN-1 exhibits balanced dual activities of less than 10 μ M on the two targets^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

An MTS assay in which the proliferation rates of HEK-Blue NOD1 cells are measured in the presence of Noditinib-1 and of the synthesized potential NOD1 inhibitor NOD-IN-1 is employed to screen these compounds for potential cytotoxicity. Cells are treated for 24 h with the compound of interest at concentrations of up to 25 μ M. Comparison of the resulting metabolic activities with that of the untreated control showed that all compounds are well tolerated by HEK-Blue NOD1 cells, since their residual metabolic activities do not fall below 80% at the maximum concentration tested^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Hazard Mater. 2020 Jul.
- Front Cell Infect Microbiol. 2020 May.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Kecek Plesec K, et al. Identification of indole scaffold-based dual inhibitors of NOD1 and NOD2. Bioorg Med Chem. 2016 Nov 1;24(21):5221-5234.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA