


## 4E2RCat

<b>Cat. No.:</b>	HY-100733												
<b>CAS No.:</b>	432499-63-3												
<b>Molecular Formula:</b>	C <sub>22</sub> H <sub>14</sub> ClNO <sub>4</sub> S <sub>2</sub>												
<b>Molecular Weight:</b>	455.93												
<b>Target:</b>	Eukaryotic Initiation Factor (eIF); Autophagy; Virus Protease												
<b>Pathway:</b>	Cell Cycle/DNA Damage; Autophagy; Anti-infection												
<b>Storage:</b>	<table border="0"> <tr> <td>Powder</td> <td>-20°C</td> <td>3 years</td> </tr> <tr> <td></td> <td>4°C</td> <td>2 years</td> </tr> <tr> <td>In solvent</td> <td>-80°C</td> <td>6 months</td> </tr> <tr> <td></td> <td>-20°C</td> <td>1 month</td> </tr> </table>	Powder	-20°C	3 years		4°C	2 years	In solvent	-80°C	6 months		-20°C	1 month
Powder	-20°C	3 years											
	4°C	2 years											
In solvent	-80°C	6 months											
	-20°C	1 month											



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 23.33 mg/mL (51.17 mM; Need ultrasonic)																					
	<table border="1"> <thead> <tr> <th rowspan="2">Solvent</th> <th rowspan="2">Mass</th> <th colspan="3">Concentration</th> </tr> <tr> <th>1 mg</th> <th>5 mg</th> <th>10 mg</th> </tr> </thead> <tbody> <tr> <td rowspan="3">Preparing Stock Solutions</td> <td>1 mM</td> <td>2.1933 mL</td> <td>10.9666 mL</td> <td>21.9332 mL</td> </tr> <tr> <td>5 mM</td> <td>0.4387 mL</td> <td>2.1933 mL</td> <td>4.3866 mL</td> </tr> <tr> <td>10 mM</td> <td>0.2193 mL</td> <td>1.0967 mL</td> <td>2.1933 mL</td> </tr> </tbody> </table>	Solvent	Mass	Concentration			1 mg	5 mg	10 mg	Preparing Stock Solutions	1 mM	2.1933 mL	10.9666 mL	21.9332 mL	5 mM	0.4387 mL	2.1933 mL	4.3866 mL	10 mM	0.2193 mL	1.0967 mL	2.1933 mL
Solvent	Mass			Concentration																		
		1 mg	5 mg	10 mg																		
Preparing Stock Solutions	1 mM	2.1933 mL	10.9666 mL	21.9332 mL																		
	5 mM	0.4387 mL	2.1933 mL	4.3866 mL																		
	10 mM	0.2193 mL	1.0967 mL	2.1933 mL																		
	Please refer to the solubility information to select the appropriate solvent.																					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.33 mg/mL (5.11 mM); Suspended solution; Need ultrasonic																					

### BIOLOGICAL ACTIVITY

<b>Description</b>	4E2RCat is an inhibitor of eIF4E-eIF4G interaction with an IC <sub>50</sub> of 13.5 μM.
<b>IC<sub>50</sub> &amp; Target</b>	IC <sub>50</sub> : 13.5 μM (eIF4E-eIF4G) <sup>[1]</sup>
<b>In Vitro</b>	4E2RCat prevents the interaction between eIF4E (the cap-binding protein) and eIF4G (a large scaffolding protein), inhibiting cap-dependent translation. It significantly decreases human coronavirus 229E (HCoV-229E) replication, reducing the percentage of infected cells and intra- and extracellular infectious virus titers. 4E2RCat inhibits cap-dependent translation in a dose-dependent manner. 4E2RCat inhibits cap-dependent FF translation but not EMCV IRES-driven Ren translation. 4E2RCat inhibits coronavirus replication in a dose- and time-dependent manner <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
<b>In Vivo</b>	4E2RCat inhibits protein synthesis in vivo and it is not a consequence of increased cell death <sup>[1]</sup> .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Cell Assay <sup>[1]</sup>

L132 cells are treated with 12.5  $\mu$ M 4E2RCat for the indicated times and are processed for annexin V/propidium iodide staining. To this end, cell medium is collected. Cells are washed with 1 mL PBS, which is collected as well, and trypsinized in 200  $\mu$ L 0.05% trypsin-EDTA. Cells are pooled with previously collected supernatants and spun for 2 min at 2,000 rpm and 4°C. The cell pellet is washed with 2 mL cold PBS, followed by another spin. After the second spin, the cell pellet is resuspended in 100  $\mu$ L annexin V binding buffer and propidium iodide added to a final concentration of 5  $\mu$ g/mL. After the addition of 5  $\mu$ L annexin V-fluorescein isothiocyanate, samples are incubated for 15 min in the dark at room temperature and diluted. Fluorescence-activated cell sorter (FACS) analyses are performed using a FACScan instrument<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- bioRxiv. May 27, 2021.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Cencic R, et al. Blocking eIF4E-eIF4G interaction as a strategy to impair coronavirus replication. J Virol. 2011 Jul;85(13):6381-9.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA