


Olcegepant

Cat. No.:	HY-10095		
CAS No.:	204697-65-4		
Molecular Formula:	C ₃₈ H ₄₇ Br ₂ N ₉ O ₅		
Molecular Weight:	869.65		
Target:	CGRP Receptor		
Pathway:	GPCR/G Protein; Neuronal Signaling		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 50 mg/mL (57.49 mM)
 H₂O : < 0.1 mg/mL (insoluble)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	1.1499 mL	5.7494 mL	11.4989 mL
	5 mM	0.2300 mL	1.1499 mL	2.2998 mL
	10 mM	0.1150 mL	0.5749 mL	1.1499 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2.5 mg/mL (2.87 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: ≥ 2.5 mg/mL (2.87 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Olcegepant (BIBN-4096) is a potent and selective non-peptide antagonist of the calcitonin gene-related peptide 1 (CGRP1) receptor with IC₅₀ of 0.03 nM and K_i of 14.4 pM for human CGRP^{[1][2]}.

IC₅₀ & Target

IC₅₀: 0.03 nM (CGRP1)^[1]
 Ki: 14.4 pM (hCGRP)^[2]

In Vitro

Olcegepant possesses higher affinity for the human CGRP receptor than the endogenous ligand CGRP and 150-fold higher affinity compared to the peptidic antagonist CGRP8-37. Olcegepant reverses CGRP-mediated vasodilation in human cerebral

vessels and inhibits neurogenic vasodilation in a surrogate animal model of migraine pathophysiology^[1]. Olcegepant (BIBN4096BS) is extremely potent at primate CGRP receptors exhibiting an affinity (K_i) for human CGRP receptors of 14.4 ± 6.3 (n=4) pM^[2]. Several lines of evidence suggest that a calcitonin-gene related peptide (CGRP) receptor antagonist may serve as a novel abortive migraine treatment. Olcegepant (BIBN4096BS) exhibits competitive antagonism at the CGRP receptor present in SK-N-MC cells. Isolated human cerebral, coronary, and omental arteries are studied with a sensitive myograph technique. CGRP induces a concentration-dependent relaxation that is antagonized by Olcegepant in a competitive manner^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Olcegepant (BIBN4096BS) in doses between 1 and 30 $\mu\text{g}/\text{kg}$ (i.v.) inhibits the effects of CGRP, released by stimulation of the trigeminal ganglion, on facial blood flow in marmoset monkeys^[2]. Pre-treatment with Olcegepant (900 $\mu\text{g}/\text{kg}$) inhibits the capsaicin-induced expression of Fos throughout the spinal trigeminal nucleus by 57%. In contrast, the expression of phosphorylated extracellular signal-regulated kinase in the trigeminal ganglion is not changed by Olcegepant pre-treatment^[4]. Olcegepant (0.3 to 0.9 mg/kg, i.v.) markedly reduces mechanical allodynia in CCI-ION rats. Olcegepant (0.6 mg/kg, i.v.) significantly reduces the number of c-Fos immunolabeled cells in spinal nucleus of the trigeminal nerve and upregulation of ATF3 transcript (a marker of neuron injury) but not that of interleukin-6 in trigeminal ganglion of CCI-ION rats^[5].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay^[2]

¹²⁵I-hCGRP is used as the radioligand. The incubation buffer contained (in mM): Tris 50, NaCl 150, MgCl₂ 5 and EDTA 1, (ethylene diamine tetra-acetic acid) pH 7.4. Membrane homogenates are incubated for 180 min at room temperature with 50 pM ¹²⁵I-hCGRP and increasing concentrations of Olcegepant (BIBN4096BS). The incubation is terminated by filtration through GF/B glass fibre filters using a cell harvester. The protein-bound radioactivity is determined in a gamma counter. The nonspecific binding is defined as radioactivity bound in the presence of 1 μM CGRP. The IC₅₀ values are obtained by non-linear regression analysis on the basis of a one binding site model^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay^[3]

Cells are washed with phosphate-buffered saline then pre-incubated with 300 μM isobutylmethylxanthine in serum-free MEM for 30 min at 37 °C α -CGRP-(S-37) or Olcegepant (BIBN4096BS) is added and the cells are incubated for 10 min before the addition of CGRP. The incubation is continued for another 15 min, then the cells are washed with PBS and processed for cAMP determination. Maximal stimulation over basal is defined by using 100 nM CGRP. Dose-response curves are generated by using Prism^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration^[5]

Rats are treated acutely with Olcegepant (0.3, 0.6, and 0.9 mg/kg, intravenously [i.v.] in a tail vein), GR-85548A (0.1 and 0.3 mg/kg subcutaneously [s.c.]), or their respective vehicle. For combined treatment, Olcegepant (0.3 mg/kg, i.v.) is administered 30 minutes before GR-85548A (0.1 mg/kg, s.c.). The doses and routes of administration are based on previous reports.

For subchronic treatment, CCI-ION and sham-operated rats are injected twice per day for 4 days (at 10 am and 6 pm) with Olcegepant (0.6 mg/kg, i.v.) or its vehicle, starting on the 15th day after ligature. A further injection of Olcegepant (0.6 mg/kg, i.v.) or vehicle is performed at 10 am the subsequent day (19th day after ligature), just before von Frey filament testing.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nat Med. 2016 Oct;22(10):1160-1169.
- J Invest Dermatol. 2019 Mar;139(3):656-664.

- Cephalalgia. 2020 Oct 15;333102420963857.
- Cephalalgia. 2020 Mar 29;333102420914913.
- Cephalalgia. 2019 Dec;39(14):1827-1837.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Rudolf K, et al. Development of human calcitonin gene-related peptide (CGRP) receptor antagonists. 1. Potent and selective small molecule CGRP antagonists. 1-[N2-[3,5-dibromo-N-[[4-(3,4-dihydro-2(1H)-oxoquinazolin-3-yl)-1-piperidinyl]carbonyl]-D-tyrosyl]-L-lysyl]-4-(4-pyridinyl)piperazine: the first CGRP antagonist for clinical trials in acute migraine. *J Med Chem.* 2005 Sep 22;48(19):5921-31.
- [2]. Doods H, et al. Pharmacological profile of BIBN4096BS, the first selective small molecule CGRP antagonist. *Br J Pharmacol.* 2000 Feb;129(3):420-3.
- [3]. Edvinsson L, et al. Effect of the CGRP receptor antagonist BIBN4096BS in human cerebral, coronary and omental arteries and in SK-N-MC cells. *Eur J Pharmacol.* 2002 Jan 2;434(1-2):49-53.
- [4]. Sixt ML, et al. Calcitonin gene-related peptide receptor antagonist Olcegepant acts in the spinal trigeminal nucleus. *Brain.* 2009 Nov;132(Pt 11):3134-41.
- [5]. Michot B, et al. Differential effects of calcitonin gene-related peptide receptor blockade by Olcegepant on mechanical allodynia induced by ligation of the infraorbital nerve vs the sciatic nerve in the rat. *Pain.* 2012 Sep;153(9):1939-48.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA