


IT1t

Cat. No.:	HY-101458
CAS No.:	864677-55-4
Molecular Formula:	C ₂₁ H ₃₄ N ₄ S ₂
Molecular Weight:	406.65
Target:	CXCR; HIV
Pathway:	GPCR/G Protein; Immunology/Inflammation; Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	IT1t is a potent CXCR4 antagonist; inhibits CXCL12/CXCR4 interaction with an IC ₅₀ of 2.1 nM.		
IC₅₀ & Target	CXCL12/CXCR4 2.1 nM (IC ₅₀)	HIV-1 (X4) 14.2 nM (IC ₅₀ , in MT-4 cells)	HIV-1 (X4) 19 nM (IC ₅₀ , in PBMCs)
In Vitro	<p>The CXCR4 is involved in chemotaxis and serves as a coreceptor for T-tropic HIV-1 viral entry and in cancer metastasis. IT1t is a small, drug-like, isothiourea derivative. IT1t shows very potent and dose-dependent inhibition of the CXCL12/CXCR4 interaction with an IC₅₀ of 2.1 nM. This calcium flux is also inhibited by IT1t with an IC₅₀ of 23.1^[1]. Strong electron density is observed for IT1t in the binding cavity of both subunits of the CXCR4 homodimer. In dimers of CXCR4 bound to IT1t, the monomers interact only at the extracellular side of helices V and VI, leaving at least a 4 Å gap between the intracellular regions, which is presumably filled by lipids. The IT1t compound and CVX15 peptide have both been characterized as competitive inhibitors of CXCL12, and many of the receptor-ligand contacts in the co-crystal structures presented are important for CXCL12 binding, including the acidic Asp187, Glu2887.39 and Asp972.63. The binding site of IT1t may point to the major anchor region for this domain^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>		
In Vivo	<p>IT1t reduces the formation of TNBC early metastases in the zebrafish xenograft model. Tumor cell invasion at the metastatic site is effectively reduced upon CXCR4 silencing (Fig. 7B), similar to the antagonist IT1t^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>		

PROTOCOL

Cell Assay^[1]	<p>Jurkat cells are incubated with serial dilutions (0.001, 0.01, 0.1, 1, 10, 100, 1000 μM) of IT1t at room temperature for two hours. Cytotoxicity of IT1t is also evaluated at 37°C over a longer period of time in MT-4 cells and PHA-stimulated PBMCs (ten day incubation) because these cell types are used in anti-HIV activity assays which last up to ten days. Cytotoxicity is evaluated microscopically and viability is assessed using the a kit^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
---------------------------------	---

CUSTOMER VALIDATION

- Nat Commun. 2021 Jul 22;12(1):4457.
- Biomed Pharmacother. 2022 Sep 27;155:113739.
- Clin Exp Pharmacol Physiol. 2024 May;51(5):e13857.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Van Hout A, et al. Comparison of cell-based assays for the identification and evaluation of competitive CXCR4 inhibitors. PLoS One. 2017 Apr 14;12(4):e0176057.
- [2]. Wu B, et al. Structures of the CXCR4 chemokine GPCR with small-molecule and cyclic peptide antagonists. Science. 2010 Nov 19;330(6007):1066-71.
- [3]. Tulotta C, et al. Inhibition of signaling between human CXCR4 and zebrafish ligands by the small molecule IT1timpairs the formation of triple-negative breast cancer early metastases in a zebrafish xenograft model. Dis Model Mech. 2016 Feb;9(2):141-53.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA