


## 9-Hydroxyellipticine hydrochloride

<b>Cat. No.:</b>	HY-101775A
<b>CAS No.:</b>	52238-35-4
<b>Molecular Formula:</b>	C <sub>17</sub> H <sub>15</sub> ClN <sub>2</sub> O
<b>Molecular Weight:</b>	298.77
<b>Target:</b>	Topoisomerase
<b>Pathway:</b>	Cell Cycle/DNA Damage
<b>Storage:</b>	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	9-Hydroxyellipticine hydrochloride is a inhibitor of Topo II and RyR. 9-Hydroxyellipticine hydrochloride exhibits antitumor, antioxidant and catecholamine-releasing activities. 9-Hydroxyellipticine hydrochloride exhibits IC <sub>50</sub> values of 1.6 μM and 1.2 μM in Hela S-3 and 293T cells, respectively <sup>[1][2][3]</sup> .	
<b>IC<sub>50</sub> &amp; Target</b>	Topoisomerase II	
<b>In Vitro</b>	9-Hydroxyellipticine (9HE) causes selective inhibition of p53 protein phosphorylation in Lewis lung carcinoma and SW480 (human colon cancer cell line) in a concentration-dependent manner from 0.1 to 100 μM <sup>[4]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
<b>In Vivo</b>	9-Hydroxyellipticine (5 or 10 mg/kg, ip) results in chromosome clumping and sister chromatid exchange in murine bone marrow cells <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	<b>Animal Model:</b>	Three- to five-month-old C57B1/6 male mice <sup>[1]</sup> .
	<b>Dosage:</b>	5 or 10 mg/kg.
	<b>Administration:</b>	IP.
	<b>Result:</b>	Resulted in chromosome clumping, chromatid aberrations, and micronuclei formation in murine bone marrow cells.

### REFERENCES

- [1]. Renault G, et al. In vivo exposure to four ellipticine derivatives with topoisomerase inhibitory activity results in chromosome clumping and sister chromatid exchange in murine bone marrow cells. *Toxicol Appl Pharmacol.* 1987 Jun 30;89(2):281-6.
- [2]. Saeki K, et al. Cardioprotective effects of 9-hydroxyellipticine on ischemia and reperfusion in isolated rat heart. *Jpn J Pharmacol.* 2002 May;89(1):21-8.
- [3]. Naoya Sato, et al. Synthesis and in vitro Antitumor Activity of 9-Hydroxyellipticine Derivatives with Glucose Conjugation via Triazolymethyl Succinate Linker. *Heterocycle*, Vol. 92, NO 4, 2016.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA