


EDO-S101

Cat. No.:	HY-101780		
CAS No.:	1236199-60-2		
Molecular Formula:	C ₁₉ H ₂₈ Cl ₂ N ₄ O ₂		
Molecular Weight:	415.36		
Target:	HDAC		
Pathway:	Cell Cycle/DNA Damage; Epigenetics		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (240.76 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	2.4076 mL	12.0378 mL	24.0755 mL
		5 mM	0.4815 mL	2.4076 mL	4.8151 mL
10 mM		0.2408 mL	1.2038 mL	2.4076 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (6.02 mM); Suspended solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	EDO-S101 (Tinostamustine) is a pan HDAC inhibitor; inhibits HDAC6, HDAC1, HDAC2 and HDAC3 with IC ₅₀ values of 9 nM, 9 nM and 25 nM, respectively ^[1] .			
IC₅₀ & Target	HDAC6 6 nM (IC ₅₀)	HDAC1 9 nM (IC ₅₀)	HDAC2 9 nM (IC ₅₀)	HDAC3 25 nM (IC ₅₀)
	HDAC10 72 nM (IC ₅₀)	HDAC8 107 nM (IC ₅₀)		
In Vitro	EDOS101 inhibits HDAC activity in rat peripheral blood mononuclear cells (PBMCs) in a cellular assay by approximately 90% one hour after dosing with 10mg/kg i.v. HDAC inhibition in PBMCs could not be increased with higher doses up to 50mg/kg. EDO-S101 triggers apoptosis and shows strong antitumor activity in HL60 and Daudi cells. Initial in vitro experiments in HL60			

cells shows an activation of the intrinsic pathway of apoptosis with cleavage of caspases 3, 9 and PARP and a marked reduction of anti-apoptotic proteins XIAP and Mcl-1^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Intracellular HDAC inhibition of EDO-S101, which occurs rapidly after dosing is at maximum activity already at the lowest dose of 10mg/kg and lasts for about 12-16 hours. Exposure to EDO-S101 causes a strong DNA repair response evidenced by activation of pH2AX and p53 in tumors taken from mice bearing subcutaneous human Burkitt's lymphoma. Tumors of BL rapidly shrink or are completely eradicated after i.v. administration of EDO-S101^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

EDO-S101 is dissolved in DMSO and added to the assay buffer solution. EDO-S101 dilutions of 5 µL of each dilution is added to 50 µL of the reaction mixture including the Fluor de Lys substrate and all of the enzymatic reactions are conducted in duplicate at 37°C for 30 minutes. After enzymatic reactions, 50 µL of 2xHDAC developer is added to each well and fluorescence intensity is measured^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Rats: The duration of HDAC inhibition is assessed in 12 female rats after receiving a single dose of either vehicle or EDO-S101 (25mg/kg). Blood samples from EDO-S101 treated rats are collected 1hr, 3hr, 6hr, 16hr and 24hr post dosing (n=2 per time point). Blood sample from vehicle treated rats (n=2) are collected 24hr post dosing^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Mehrling T, et al. The Alkylating-HDAC Inhibition Fusion Principle: Taking Chemotherapy to the Next Level with the First in Class Molecule EDO-S101. Anticancer Agents Med Chem. 2016;16(1):20-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA