


## BQCA

<b>Cat. No.:</b>	HY-101858		
<b>CAS No.:</b>	338747-41-4		
<b>Molecular Formula:</b>	C <sub>18</sub> H <sub>15</sub> NO <sub>4</sub>		
<b>Molecular Weight:</b>	309.32		
<b>Target:</b>	mAChR		
<b>Pathway:</b>	GPCR/G Protein; Neuronal Signaling		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



## SOLVENT & SOLUBILITY

### In Vitro

DMSO : 5.45 mg/mL (17.62 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	3.2329 mL	16.1645 mL	32.3290 mL
	5 mM	0.6466 mL	3.2329 mL	6.4658 mL
	10 mM	0.3233 mL	1.6164 mL	3.2329 mL

Please refer to the solubility information to select the appropriate solvent.

## BIOLOGICAL ACTIVITY

### Description

BQCA a highly selective allosteric modulator of the M1 mAChR.

### In Vitro

BQCA reduces the concentration of ACh required to activate M1 up to 129-fold with an inflection point value of 845 nM. No potentiation, agonism, or antagonism activity on other mAChRs is observed up to 100 μM<sup>[1]</sup>. BQCA increases M1 receptor affinity for acetylcholine. The activation of the M1 receptor by BQCA induces a robust inward current and increases spontaneous excitatory postsynaptic currents in medial prefrontal cortex (mPFC) pyramidal cells<sup>[2]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### In Vivo

BQCA requires M1 to promote inositol phosphate turnover in primary neurons and to increase c-fos and arc RNA expression and ERK phosphorylation in the brain. BQCA reverses scopolamine-induced memory deficits in contextual fear conditioning, increases blood flow to the cerebral cortex, and increases wakefulness while reducing delta sleep. BQCA induces β-arrestin recruitment to M1, suggesting a role for this signal transduction mechanism in the cholinergic modulation of memory<sup>[1]</sup>. BQCA increases firing of mPFC pyramidal cells in vivo. BQCA also restores discrimination reversal learning in a transgenic mouse model of Alzheimer's disease<sup>[2]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Kinase Assay <sup>[1]</sup>

Competition binding reactions used 25 µg human M1 CHO membrane protein, BQCA or vehicle, and 0.15 nM [<sup>3</sup>H]NMS in 96-well deep-well plates. Binding reactions (30 °C for 2-3 h) are terminated by rapid filtration. Nonspecific binding is determined by adding 10 µM atropine. Filter plates are washed 4× with ice-cold 20 mM HEPES, 100 mM NaCl, and 5 mM MgCl<sub>2</sub>, pH 7.4 using a 96-well harvester. Plates are dried and radioactivity counted with a microplate scintillation counter<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Animal Administration <sup>[1][2]</sup>

Rats: Male Sprague-Dawley rats weighing 225-250 g, are injected i.p. with the micro-suspension (containing 10% tween 80) of BQCA at the dose of 10 mg/kg. The blood and whole brain tissue samples are collected at 0.5, 1, 2, 4 and 8 h. Blood samples are collected through cardiac puncture in EDTA vacutainer tubes. The plasma is separated by centrifugation and stored at -80°C until analysis. The animals are decapitated and the whole brain tissue are removed and immediately frozen on dry ice<sup>[2]</sup>.

Mice: Mice are dosed I.P. with BQCA in 5% beta-cyclodextrin and/or 0.3 mg/kg scopolamine in 0.9% saline 30 min before placement into a chamber for 2 min before 2 tone-footshock pairings (3 kHz, 85 dB tone for 30 s co-terminated with a 0.5 mA, 1 s shock) 2 min apart. Mice are removed to their home cage 30 s after the last pairing. Twenty-four hours later mice are placed into the same chamber and freezing is measured by Video Freeze<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

- [1]. Ma L, et al. Selective activation of the M1 muscarinic acetylcholine receptor achieved by allosteric potentiation. Proc Natl Acad Sci U S A. 2009 Sep 15;106(37):15950-5.
- [2]. Shirey JK, et al. A selective allosteric potentiator of the M1 muscarinic acetylcholine receptor increases activity of medial prefrontal cortical neurons and restores impairments in reversal learning. J Neurosci. 2009 Nov 11;29(45):14271-86.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA