


Sepantronium bromide

Cat. No.:	HY-10194		
CAS No.:	781661-94-7		
Molecular Formula:	C ₂₀ H ₁₉ BrN ₄ O ₃		
Molecular Weight:	443.29		
Target:	Survivin; Autophagy		
Pathway:	Apoptosis; Autophagy		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (225.59 mM; Need ultrasonic)
 DMSO : 50 mg/mL (112.79 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.2559 mL	11.2793 mL	22.5586 mL
	5 mM	0.4512 mL	2.2559 mL	4.5117 mL
	10 mM	0.2256 mL	1.1279 mL	2.2559 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 50 mg/mL (112.79 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2 mg/mL (4.51 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: 2 mg/mL (4.51 mM); Clear solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

Sepantronium bromide (YM-155) is a survivin inhibitor with an IC₅₀ of 0.54 nM^[1].

IC₅₀ & Target

IC₅₀: 0.54 nM (Survivin)^[1]

In Vitro

Sepantronium bromide (YM155; 30 μM) is not sensitive to survivin gene promoter-driven luciferase reporter activity. Sepantronium bromide shows significant suppression on endogenous survivin expression in PC-3 and PPC-1 human HRPC

cells with deficient p53 via transcriptional inhibition of the survivin gene promoter. Sepantronium bromide (100 nM) does not affect protein expression of c-IAP2, XIAP, Bcl-2, Bcl-xL, Bad, α -actin, and β -tubulin. Sepantronium bromide potently inhibits human cancer cell lines (mutated or truncated p53) such as PC-3, PPC-1, DU145, TSU-Pr1, 22Rv1, SK-MEL-5 and A375 with IC₅₀s ranging from 2.3 to 11 nM, respectively^[1].

Sepantronium bromide (YM155) result in an increase in sensitivity of NSCLC cells to γ -radiation. Sepantronium bromide combined with γ -radiation increases both the number of apoptotic cells and the activity of caspase-3. In addition, Sepantronium bromide delays the repair of radiation-induced double-strand breaks in nuclear DNA^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Sepantronium bromide (YM155; 3 and 10 mg/kg) inhibits the tumor growth in PC-3 xenografts, without obvious body weight loss and blood cell count decrease. Sepantronium bromide is highly distributed to tumor tissue in vivo. Sepantronium bromide shows 80% TGI at a dose of 5 mg/kg in PC-3 orthotopic xenografts^[1].

Sepantronium bromide (YM155) in combination with γ -radiation shows potent antitumor activity against H460 or Calu6 xenografts in nude mice^[2].

In this orthotopic renal and metastatic lung tumors models, Sepantronium bromide (YM-155) and IL-2 additively decreases tumor weight, lung metastasis, and luciferin-stained tumor images^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

The antiproliferative activity of Sepantronium bromide is measured. After treatment with Sepantronium bromide for 48 h, the cell count is determined by sulforhodamine B assay. The GI₅₀ value is calculated by logistic analysis, which is the drug concentration resulting in a 50% reduction in the net protein increase (as measured by sulforhodamine B staining) in control cells during the drug incubation. The assay is done in triplicate, and the mean GI₅₀ value is obtained from the results of four independent assays.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Five-week-old male nude mice (BALB/c nu/nu) are used for the assay. PC-3 cells (2×10^6 - 3×10^6) are injected into the flanks of the mice and allowed to reach a tumor volume of $> 100 \text{ mm}^3$ in tumor volume ($\text{length} \times \text{width}^2 \times 0.5$). Sepantronium bromide is s.c. administered as a 3-day continuous infusion per week for 2 weeks using an implanted micro-osmotic pump or i.v. administered five times a week for 2 weeks. The percentage of tumor growth inhibition 14 days after initial Sepantronium bromide administration is calculated for each group using the following formula: $\text{MTV} = 100 \times \{1 - [(\text{MTV of the treated group on day 14}) - (\text{MTV of the treated group on day 0})] / [(\text{MTV of the control group on day 14}) - (\text{MTV of the control group on day 0})]\}$, where MTV is mean tumor volume. For both the frozen tumors and plasma samples, survivin expression levels are analyzed by Western blotting and Sepantronium bromide concentration by high-performance liquid chromatography/triple quadrupole mass spectrometry (LC/MS/MS) using validated methods.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cancer Lett. 2018 Jul 1;425:54-64.
- Cancers. 2019 Oct 14;11(10):1550.
- Cancers. 2019 Jul 5;11(7):947.
- Stem Cell Res Ther. 2020 Jun 10;11(1):229.
- Nutrients. 2018 Mar 15;10(3). pii: E353.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Nakahara T, et al. YM155, a novel small-molecule survivin suppressant, induces regression of established human hormone-refractory prostate tumor xenografts. *Cancer Res.* 2007 Sep 1;67(17):8014-21.
- [2]. Iisa T, et al. Radiosensitizing effect of YM155, a novel small-molecule survivin suppressant, in non-small cell lung cancer cell lines. *Clin Cancer Res.* 2008 Oct 15;14(20):6496-504.
- [3]. Guo K, et al. A combination of YM-155, a small molecule survivin inhibitor, and IL-2 potently suppresses renal cell carcinoma in murine model. *Oncotarget.* 2015 Aug 28;6(25):21137-47.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA