


PD0325901

Cat. No.:	HY-10254		
CAS No.:	391210-10-9		
Molecular Formula:	C ₁₆ H ₁₄ F ₃ IN ₂ O ₄		
Molecular Weight:	482.19		
Target:	MEK; Autophagy; Apoptosis		
Pathway:	MAPK/ERK Pathway; Autophagy; Apoptosis		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 56 mg/mL (116.14 mM)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.0739 mL	10.3694 mL	20.7387 mL
	5 mM	0.4148 mL	2.0739 mL	4.1477 mL
	10 mM	0.2074 mL	1.0369 mL	2.0739 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (5.18 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (5.18 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (5.18 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

PD0325901 (PD325901) is an orally active, selective and non-ATP-competitive MEK inhibitor with an IC₅₀ of 0.33 nM. PD0325901 exhibits a K_i^{APP} of 1 nM against activated MEK1 and MEK2. PD0325901 suppresses the expression of p-ERK1/2 and induces apoptosis. PD0325901 has anti-cancer activity for a broad spectrum of human tumor xenografts^{[1][2][3]}.

IC₅₀ & Target

MEK1 1 nM (K _i)	MEK2 1 nM (K _i)	MEK 0.33 nM (IC ₅₀)
--------------------------------	--------------------------------	------------------------------------

In Vitro	<p>PD0325901 (PD325901; 0.0064, 0.032, 0.16, 0.8, 4, 20, 100 nM; for 2 days) inhibits the growth of Papillary thyroid carcinomas (PTC) cell lines (TPC-1 cells and K2 cells) with GC₅₀ of 11 nM and 6.3 nM, respectively^[3].</p> <p>PD0325901 (100 nmol/L; for 4 days) induces apoptosis in K2 cells (top) or TPC-1 cells^[3].</p> <p>PD0325901 (0.1, 1, 10, 100, 1000 nM; for 1 hour) suppresses the expression of p-ERK1/2 in K2 cells (top) or TPC-1 cells^[3].</p> <p>PD0325901 prevents the growth of melanoma cell lines. PD0325901 significantly prevents the the growth of PTC cells harboring a BRAF mutation at very low concentration (10 nM)^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>PD0325901 (PD325901; 25 mg/kg, p.o.) inhibits phosphorylation of ERK by more than 50% at 24 hours post-dosing.</p> <p>PD0325901 (25 mg/kg/day; po) produces a 70% incidence of complete tumor responses (C26 model)^[2].</p> <p>PD0325901 (20-25 mg/kg/day; oral gavage; for 3 weeks (5 consecutive days/week)) suppresses tumor growth completely in mice inoculated with PTC cells carrying a BRAF mutation (K2) and significantly decreased tumor growth in mice inoculated with PTC cells carrying the RET/PTC1 rearrangement (TPC-1) in athymic Ncr-nu/nu mice at ages 6 to 8 weeks^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

PROTOCOL

Kinase Assay ^[1]	<p>Incorporation of ³²P into myelin basic protein (MBP) is assayed in the presence of a glutathione S-transferase fusion protein containing p44MAP kinase (GST-MAPK) and a glutathione S-transferase protein containing p45MEK (GST-MEK). The assay solution contained 20 mM HEPES, pH 7.4, 10 mM MgCl₂, 1 mM MnCl₂, 1 mM EGTA, 50 mM [gamma-³²P]ATP, 10 mg GST-MEK, 0.5 mg GST-MAPK and 40 mg MBP in a final volume of 100 mL. Reactions are stopped after 20 minutes by addition of trichloroacetic acid and filtered through a GF/C filter mat. ³²P retained on the filter mat is determined using a 1205 Betaplate. PD0325901 is assessed at various dose ranges in order to determine dose response curves.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Cell Assay ^[2]	<p>PTC cells (1×10⁴) are seeded in 24-well plates with 1 mL of medium for 4 days in a 37°C incubator. MEK inhibitor PD0325901 at varying concentrations is added to the cells in triplicate on day 0. MTT dissolved in 0.8% NaCl solution at 5 mg/mL is added to each well (0.2 mL) on day 2 to test GC₅₀ or every day for cell growth curves. The cells are incubated at 37°C for 3 hours with MTT. The liquid is then aspirated from the wells and discarded. Stained cells are dissolved in 0.5 mL of DMSO and their absorption at 570 nm is measured using a Synergy HT multidetection microplate reader. For GC₅₀, cell growth is calculated as $100 \times (T - T_0) / (C - T_0)$, where T is the optical density of the wells treated with inhibitors after a 48-hour period, T₀ is the optical density at time zero, and C is the control optical density with DMSO only.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Animal Administration ^[2]	<p>Mice (10-14 per group) are anesthetized s.c. with a cocktail. K2 and TPC-1 cells stably infected with a retrovirus expressing luciferase (5×10⁵ cells in 5 μL RPMI1640 medium) are inoculated into the thyroid gland, and the mice are monitored weekly for tumor growth by Xenogen using Living Image 3.0 software. One week after inoculation, PD0325901 is dissolved in 80 mM citric buffer (pH 7) by sonication and given to mice daily by oral gavage (20-25 mg/kg) for 3 weeks (5 consecutive days/week). Mice are sacrificed only due to tumor burden or loss of 20% of body weight. Tumor sizes are measured with calipers and tumor volume (V) is calculated by the formula (V=length×width×depth). Control mice are given 80 mM citric buffer (pH 7) alone. All in vivo experiments are done at least twice.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

CUSTOMER VALIDATION

- Nat Biomed Eng. 2018 Aug;2(8):578-588.
- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Biomaterials. 2018 Oct;180:12-23.

- Neuro Oncol. 2019 Mar 18;21(4):486-497.
- Clin Cancer Res. 2020 Apr 15;26(8):2011-2021.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Barrett SD, et al. The discovery of the benzhydroxamate MEK inhibitors CI-1040 and PD 0325901. Bioorg Med Chem Lett. 2008 Dec 15;18(24):6501-4.
- [2]. Henderson YC, et al. MEK inhibitor PD0325901 significantly reduces the growth of papillary thyroid carcinoma cells in vitro and in vivo. Mol Cancer Ther. 2010 Jul;9(7):1968-76.
- [3]. Judith S. Sebolt-Leopold, et al. The biological profile of PD 0325901: A second generation analog of CI-1040 with improved pharmaceutical potential
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA