


## Ensartinib hydrochloride

|                    |  |       |          |
|--------------------|--|-------|----------|
| Cat. No.:          | HY-103714A   |       |          |
| CAS No.:           | 2137030-98-7   |       |          |
| Molecular Formula: | C <sub>26</sub> H <sub>29</sub> Cl <sub>4</sub> FN <sub>6</sub> O <sub>3</sub> |       |          |
| Molecular Weight:  | 634.36   |       |          |
| Target:            | ALK; c-Met/HGFR  |       |          |
| Pathway:           | Protein Tyrosine Kinase/RTK  |       |          |
| Storage:           | Powder   | -20°C | 3 years  |
|                    |  | 4°C   | 2 years  |
|                    | In solvent   | -80°C | 6 months |
|                    |  | -20°C | 1 month  |



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 35.71 mg/mL (56.29 mM; Need ultrasonic)  
 H<sub>2</sub>O : 33.33 mg/mL (52.54 mM; Need ultrasonic)

| Preparing Stock Solutions | Solvent Concentration | Mass      |           |            |
|---------------------------|-----------------------|-----------|-----------|------------|
|                           |                       | 1 mg      | 5 mg      | 10 mg      |
|                           | 1 mM                  | 1.5764 mL | 7.8820 mL | 15.7639 mL |
|                           | 5 mM                  | 0.3153 mL | 1.5764 mL | 3.1528 mL  |
|                           | 10 mM                 | 0.1576 mL | 0.7882 mL | 1.5764 mL  |

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.08 mg/mL (3.28 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.08 mg/mL (3.28 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 2.08 mg/mL (3.28 mM); Clear solution
- Add each solvent one by one: PBS  
Solubility: 14.29 mg/mL (22.53 mM); Clear solution; Need ultrasonic

### BIOLOGICAL ACTIVITY

#### Description

Ensartinib hydrochloride (X-396 hydrochloride) is a potent and dual ALK/MET inhibitor with IC<sub>50</sub>s of <0.4 nM and 0.74 nM, respectively.

#### IC<sub>50</sub> & Target

IC<sub>50</sub>: <0.4 nM (ALK), 0.74 nM (MET)<sup>[1]</sup>

|                 |  |
|-----------------|--|
| <b>In Vitro</b> | <p>Ensartinib (X-396) is a potent and dual ALK/MET inhibitor with IC<sub>50</sub>s of &lt;0.4 nM and 0.74 nM, respectively. Ensartinib is potent in H3122 lung cancer cells harboring EML4-ALK E13;A20 (IC<sub>50</sub>: 15 nM). Ensartinib is also potent in H2228 lung cancer cells harboring EML4-ALK E6a/b; A20 (IC<sub>50</sub>: 45 nM). Furthermore, X-376 is potent in SUDHL-1 lymphoma cells harboring NPM-ALK (IC<sub>50</sub>: 9 nM)<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>   |
| <b>In Vivo</b>  | <p>Ensartinib (X-396) shows substantial bioavailability and moderate half-lives in vivo. Nude mice harboring H3122 xenografts are treated with Ensartinib at 25 mg/kg bid. Ensartinib significantly delays the growth of tumors compared to vehicle alone. In the xenograft experiments, Ensartinib appears well-tolerated in vivo. Mouse weight is unaffected by Ensartinib treatment. Drug-treated mice appear healthy and do not display any signs of compound related toxicity. To further assess potential side effects of Ensartinib, additional systemic toxicity and toxico-kinetic studies are performed in Sprague Dawley (SD) rats. Following 10 days of repeated oral administration of Ensartinib at 20, 40, 80 mg/kg in SD rats, all animals survive to study termination. The no significant toxicity (NST) levels are determined to be 80 mg/kg for Ensartinib. At NST levels, Ensartinib achieves an AUC of 66 μM×hr and a C<sub>max</sub> of 7.19 μM<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> |

## PROTOCOL

|   |  |
|---|--|
| <b>Cell Assay</b> <sup>[1]</sup>            | <p>For viability experiments, cells are seeded in 96-well plates at 25%-33% confluency and exposed to drugs. The human lung adenocarcinoma cell lines H3122 and H2228 are treated with Ensartinib (10, 30, 100, 300 and 1000 nM). SUDHL-1 lymphoma cells are treated with Ensartinib (5, 10, 30, 100 and 300 nM). SY5Y neuroblastoma cells are treated with Ensartinib (30, 100, 300 and 1000 nM). At 72 hours post Ensartinib addition, Cell Titer Blue Reagent is added and fluorescence is measured on a Spectramax spectrophotometer. All experimental points are set up in hexuplicate replicates and are performed at least two independent times. IC<sub>50</sub>s are calculated using GraphPad Prism version 5 for Windows. The curves are fit using a nonlinear regression model with a log (inhibitor) vs. response formula<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> |
| <b>Animal Administration</b> <sup>[1]</sup> | <p>Mice<sup>[1]</sup></p> <p>Nude mice (<i>nu/nu</i>) are injected with H3122 cells. Once tumors reach an average volume of 450 mm<sup>3</sup>, a total of 27 athymic mice harboring H3122 tumors are randomized and dosed via oral gavage with 25 mg/kg Ensartinib or the control vehicle. Two, five, and fifteen hours after the single treatment (3 tumors/timepoint/group), mice are sacrificed and serum is collected for assessment of drug concentration using an LC-MS based bioanalytical method<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>  |

## CUSTOMER VALIDATION

- Cancers. 2020 Mar 28;12(4):813.
- Drug Des Dev Ther. 2020 Nov 30;14:5259-5273.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Lovly CM, et al. Insights into ALK-driven cancers revealed through development of novel ALK tyrosine kinaseinhibitors. Cancer Res. 2011 Jul 15;71(14):4920-31.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA