
## RedChemExpress

## Product Data Sheet

## Regorafenib N-oxide and N-desmethyl (M5)-<sup>13</sup>C<sub>6</sub>

| Cat. No.:          | HY-108226S1   |           |          |
|--------------------|---|-----------|----------|
| Molecular Formula: | C <sub>14</sub> <sup>13</sup> C <sub>6</sub> H <sub>13</sub> ClF <sub>4</sub> N <sub>4</sub> O <sub>4</sub> |           |          |
| Molecular Weight:  | 490.74  |           |          |
| Target:            | Isotope-Lal   | beled Cor | npounds  |
| Pathway:           | Others  |           |          |
| Storage:           | Powder  | -20°C     | 3 years  |
|                    |   | 4°C       | 2 years  |
|                    | In solvent  | -80°C     | 6 months |
|                    |   | -20°C     | 1 month  |
|                    |   |           |          |



| BIOLOGICAL ACTIVITY |   |  |  |
|---------------------|---|--|--|
| BIOLOGICAL ACTIVITY |   |  |  |
| Description         | Regorafenib N-oxide and N-desmethyl (M5)- $^{13}C_6$ is the $^{13}$ C-labeled Regorafenib N-oxide and N-desmethyl (M5)[1].  |  |  |
| In Vitro            | Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |  |  |

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-223.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA