

Simethicone

Cat. No.:	HY-109519
CAS No.:	8050-81-5
Target:	Others
Pathway:	Others
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)

Simethicone

SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (Need ultrasonic)
In Vivo	<p>1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (Infinity mM); Clear solution</p> <p>2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (Infinity mM); Clear solution</p> <p>3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (Infinity mM); Clear solution</p>

BIOLOGICAL ACTIVITY

Description	Simethicone is an orally active defoamer. Simethicone reduces the surface tension of air bubbles in the gastrointestinal tract, causing them to be expelled by vomiting, exhalation or absorption into the bloodstream. Simethicone has potential applications in flatulence and colic ^[1] .	
In Vivo	Simethicone (200 mg/kg; p.o.; single dose) has gastrointestinal regulation function in rats ^[2] . Simethicone (20 mg/kg; p.o.; single dose) shows jejunal regulation in rabbits ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	Female adult Wistar rats (200–250 g) ^[2] .
	Dosage:	50, 100 and 200 mg/kg.
	Administration:	Oral gavage; single dose.
	Result:	Improved colon permeability and hypersensitivity (200 mg/kg).
	Animal Model:	Adult New Zealand White (NZW) rabbits ^[3] .
	Dosage:	20 mg/kg.

Administration:	Oral gavage; single dose.
Result:	Improved image quality scores for the jejunum without affecting left kidney, right kidney and gallbladder.

REFERENCES

- [1]. Voepel-Lewis TD, et al. Evaluation of simethicone for the treatment of postoperative abdominal discomfort in infants. J Clin Anesth. 1998 Mar;10(2):91-4.
- [2]. Bueno L, et al. Influence of simethicone and alverine on stress-induced alterations of colonic permeability and sensitivity in rats: beneficial effect of their association. J Pharm Pharmacol. 2013 Apr;65(4):567-73.
- [3]. da Silva KG, et al. Influence of simethicone and fasting on the quality of abdominal ultrasonography in New Zealand White rabbits. Acta Vet Scand. 2017 Jul 17;59(1):48.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA