


D4-abiraterone

Cat. No.:	HY-109619
CAS No.:	154229-21-7
Molecular Formula:	C ₂₄ H ₂₉ NO
Molecular Weight:	347.49
Target:	Androgen Receptor; Cytochrome P450
Pathway:	Others; Metabolic Enzyme/Protease
Storage:	4°C, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (143.89 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
		Concentration				
		1 mM		2.8778 mL	14.3889 mL	28.7778 mL
		5 mM		0.5756 mL	2.8778 mL	5.7556 mL
10 mM		0.2878 mL	1.4389 mL	2.8778 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.19 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.19 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	D4-abiraterone is a major metabolite of abiraterone. D4-abiraterone is an inhibitor of CYP17A1, 3β-hydroxysteroid dehydrogenase (3βHSD) and steroid-5α-reductase (SRD5A) and also an antagonist of androgen receptor.
IC₅₀ & Target	CYP17A1, 3βHSD, SRD5A, androgen receptor ^[1]
In Vitro	D4-abiraterone (D4A) (10 mM) nearly completely blocks conversion from D4-androstenedione (AD) to 5α-androstenedione and other 5α-reduced androgens. The affinity of D4-abiraterone for mutant (expressed in LNCaP, half-maximum inhibitory concentration (IC ₅₀ =5.3 nM)) and wild type (expressed in LAPC4, IC ₅₀ =7.9 nM) androgen receptor (AR) is greater than that of abiraterone (Abi) (IC ₅₀ =418 and >500 nM, respectively). Compare with Abi, D4-abiraterone clearly better suppresses PSA, TMPRSS2 and FKBP5 expression in LNCaP, LAPC4 and C4-2 cell lines. D4-abiraterone also inhibits AR target gene expression in a dose-dependent manner ^[1] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

D4-abiraterone (D4A) is tenfold more potent than abiraterone (Abi) in blocking conversion from dehydroepiandrosterone (DHEA) by 3 β -hydroxysteroid dehydrogenase (3 β HSD) to D4-androstenedione (AD) in LNCaP and VCaP xenografts. 0.1 μ M D4-abiraterone is equivalent to 1 μ M Abi for blocking AD accumulation at 48 h in both LNCaP and VCaP xenografts. Progression is significantly delayed in the D4-abiraterone group compare with the Abi acetate group (P=0.011). D4-abiraterone treatment increases progression-free survival compare with Abi acetate^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

To test D4-abiraterone (D4A) as an inhibitor of 3 β HSD, enzyme assays are performed. Briefly, incubations are prepared with recombinant human 3 β HSD1 or 3 β HSD2 (in yeast microsomes, 45 or 2.5 μ g protein per incubation, respectively), D4-abiraterone (5 to 20 μ M) or ethanol vehicle in 0.5 mL of potassium phosphate buffer (pH 7.4). After a pre-incubation at 37°C for 1 to 3 min, NAD⁺ (1 mM) is added, and the incubation is conducted at 37°C for 20 min. The reaction is stopped by addition of 1 mL ethyl acetate:isooctane (1:1), and the steroids are then extracted into the organic phase and dried. The steroids in the dried extracts are resolved by HPLC and quantitated by in-line scintillation counting^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay ^[1]

Cells are cultured in serum-free medium for 48 h and then treated with the indicated concentrations of D4-abiraterone (D4A) for 30 min. Cells are washed with 1 \times PBS four times and 0.9% NaCl solution twice before lysis with RIPA buffer. Intracellular radioactivity is measured with a liquid scintillation counter and normalized to the protein concentration as detected with a Multilabel counter^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Male NSG mice, 6 to 8 weeks of age are used in this study. Mice are surgically orchiectomized and implanted with a 5 mg 90-day sustained-release dehydroepiandrosterone (DHEA) pellet to mimic castration-resistant prostate cancer (CRPC) in the context of human adrenal physiology. Two days later, 10⁷ VCaP or C4-2 cells are injected subcutaneously with matrigel. Once tumours reach 300mm³, mice are arbitrarily (but not strictly randomized) assigned to vehicle (n=9 or 10 mice for VCaP and C4-2 respectively), D4-abiraterone (D4A) (n=10 mice for both cell lines) treatment groups. D4-abiraterone (0.5 mmol per kg per day in 0.1 mL 5% benzyl alcohol and 95% safflower oil solution) is administered via 5 mL per kg intraperitoneal injection every day for up to 15 days. Control groups are administered 0.1 mL 5% benzyl alcohol and 95% safflower oil solution via intraperitoneal injection every day. Tumour volume is measured daily, and time to increase in tumour volume by 20% is determined. Mice are killed at treatment day 15 or when the tumour size is twofold greater than baseline^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Li Z, et al. Conversion of abiraterone to D4A drives anti-tumour activity in prostate cancer. Nature. 2015 Jul 16;523(7560):347-51.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA