


NIK SMI1

Cat. No.:	HY-112433		
CAS No.:	1660114-31-7		
Molecular Formula:	C ₂₀ H ₁₉ N ₃ O ₄		
Molecular Weight:	365.38		
Target:	NF-κB		
Pathway:	NF-κB		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	1 year
		-20°C	6 months



SOLVENT & SOLUBILITY

In Vitro

DMSO : 125 mg/mL (342.11 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent		1 mg	5 mg	10 mg
	Concentration	Mass			
1 mM			2.7369 mL	13.6844 mL	27.3688 mL
5 mM			0.5474 mL	2.7369 mL	5.4738 mL
10 mM			0.2737 mL	1.3684 mL	2.7369 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 5% DMSO >> 40% PEG300 >> 5% Tween-80 >> 50% saline
Solubility: ≥ 2.5 mg/mL (6.84 mM); Clear solution
- Add each solvent one by one: 5% DMSO >> 95% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (6.84 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (5.69 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.08 mg/mL (5.69 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.08 mg/mL (5.69 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

NIK SMI1 is a potent, selective NF-κB inducing kinase (NIK) inhibitor, which inhibits NIK-catalyzed hydrolysis of ATP to ADP with IC₅₀ of 0.23±0.17 nM. NIK SMI1 is a click chemistry reagent, it contains an Alkyne group and can undergo copper-catalyzed azide-alkyne cycloaddition (CuAAC) with molecules containing Azide groups.

IC₅₀ & Target	NIK ^[1]
In Vitro	<p>NIK SMI1 (Compound 4f) inhibits NIK-catalyzed hydrolysis of ATP to ADP (fluorescence polarization, FP) with an IC₅₀ of 0.23±0.17 nM. NIK SMI1 inhibits the expression of NIK SMI1 response element-regulated firefly luciferase reporter gene in HEK293 cells with an IC₅₀ of 34±6 nM. Consistent with expectations for a NIK inhibitor, NIK SMI1 is shown to inhibit nuclear translocation of p52 (RelB) (IC₅₀=70 nM). NIK SMI1 inhibits BAFF-induced B cell (mouse) survival in vitro with an IC₅₀ of 373±64 nM^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>C57BL/6 mice are treated twice daily for 7 days with orally administered NIK SMI1 or with three injections of recombinant BAFF receptor fusion protein (Br3- mIgG2a) over the course of the 7-day experiment as a positive control. The nonlinearity of exposure relative to dose between 100 and 200 mg/kg is a result of saturation of clearance mechanisms. The pharmacology of NIK SMI1 is examined in SD rat, CD-1 mouse, beagle, and cynomolgous monkey with 20, 32, 18, and 7.8 mL/kg per min, respectively. Volume of distribution (Vd, L/kg) is 1.35, 1.58, 0.778, and 1.39, respectively^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

PROTOCOL

Cell Assay ^[1]	<p>Human B cells are re-suspended in RPMI with 10% FBS for the proliferation assays and 2.5% FBS for the survival assays. Mouse B cells are plated in Co-star 96-well plates at either 50,000 cells/well for the survival assays or at 150,000 cells/well for the proliferation assays. Compounds (e.g., NIK SMI1) diluted in DMSO (final DMSO assay concentration=0.1%) are added to the cells. The cells are incubated with NIK SMI1 for one hour at 37°C. Stimulus is then added to the plates and survival or proliferation is measured after four days. For the proliferation assays, cells are treated with either Anti-IgM (20 µg/mL) or rhCD40L (10 µg/mL) or anti-mouse CD40 (100 ng/mL). For the BAFF survival assay, cells are treated with human or mouse rBAFF at 10 ng/mL followed by Cell Titer Glo to measure survival on day four^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Animal Administration ^[1]	<p>Mice^[1]</p> <p>Age-matched C57BL/6 mice are used. Only female mice are used in these experiments. The single oral doses of NIK SMI1 are 10, 20, 60, 100, and 200 mg/kg. For PO dosing, animals are manually restrained, then dosed via oral gavage using an appropriately sized gavage needle. Animals are monitored for any signs of aspiration or distress-respiratory abnormalities, lethargy, pale extremities, etc. For sample collection, 3 mice per group are bled a total of 8 times via tail prick using a 27 G needle (lateral tail vein). 10 µL of blood is collected at each timepoint and deposited into a pre-filled costar cluster tube containing 40 µL of 1.7 mg/mL EDTA/water, the tube is capped, vortexed for 5 seconds, then stored on dry ice. Samples are transferred to a -80°C freezer for storage^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

CUSTOMER VALIDATION

- Nat Immunol. 2020 May;21(5):535-545.
- Sci Immunol. 2022 Aug 12;7(74):eabn3800.
- Mol Neurobiol. 2021 Jan 13.
- J Immunol Res. 2020 Jul 31;2020:1859260.
- Research Square Print. 2022 Jun.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Blaquiere N, et al. Scaffold-Hopping Approach To Discover Potent, Selective, and Efficacious Inhibitors of NF- κ B Inducing Kinase. J Med Chem. 2018 Aug 9;61(15):6801-6813.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA