


12-HETE

Cat. No.:	HY-113439
CAS No.:	71030-37-0
Molecular Formula:	C ₂₀ H ₃₂ O ₃
Molecular Weight:	320.47
Target:	Apoptosis
Pathway:	Apoptosis
Storage:	Solution, -20°C, 2 years



BIOLOGICAL ACTIVITY

Description	<p>12-HETE, a major metabolic product of arachidonic acid using 12-LOX catalysis, inhibits cell apoptosis in a dose-dependent manner. 12-HETE promotes the activation and nuclear translocation of NF-κB through the integrin-linked kinase (ILK) pathway^[1]. 12-HETE has both anti-thrombotic and pro-thrombotic effects^[2]. 12-HETE is a neuromodulator^[3].</p>												
In Vitro	<p>12-HETE participates in the inhibition of cell apoptosis by activating the ILK/NF-κB pathway, implying an important underlying mechanism that promotes the survival of ovarian cancer cells. 12-HETE facilitates cell survival by activating the integrin-linked kinase/NF-κB pathway in ovarian cancer. 12-HETE protects against cell apoptosis in ovarian cancer cells in a concentration-dependent manner. 12-HETE (1 μM) significantly decreases the activation of caspase-3 induced by serum deprivation (SD). 12-HETE represses the increased activity of caspase-3 induced by SD in a concentration-dependent manner, with an IC₅₀ value of 1.13 μM^[1].</p> <p>12-HETE (1 μM) facilitates the activation and nuclear translocation of NF-κB via ILK in ovarian cancer cells^[1].</p> <p>12-HETE inhibits insulin secretion, reduces metabolic activity and induces cell death in human islets. 12-HETE increases bovine platelet aggregation induced by thrombin and inhibits prostaglandin E1-induced elevation of intracellular cAMP levels. 12-HETE inhibits washed platelet (WP) aggregation^[2].</p> <p>The neuronal effects of 12-HETE include attenuation of calcium influx and glutamate release as well as inhibition of AMPA receptor (AMPA-R) activation^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Viability Assay^[1]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>Ovarian cancer OVCAR-3 and SKOV3 cells</td> </tr> <tr> <td>Concentration:</td> <td>0, 0.2, 0.5, and 1 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>0, 24, 48, 72, and 96 hours</td> </tr> <tr> <td>Result:</td> <td>Inhibited the decrease in cell viability induced by SD in a dose-dependent manner. 1 μM 12-HETE treatment significantly mitigated the decrease in cell viability under conditions of SD.</td> </tr> </table> <p>Western Blot Analysis^[1]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>Ovarian cancer OVCAR-3 and SKOV3 cells</td> </tr> <tr> <td>Concentration:</td> <td>1 μM</td> </tr> </table>	Cell Line:	Ovarian cancer OVCAR-3 and SKOV3 cells	Concentration:	0, 0.2, 0.5, and 1 μM	Incubation Time:	0, 24, 48, 72, and 96 hours	Result:	Inhibited the decrease in cell viability induced by SD in a dose-dependent manner. 1 μM 12-HETE treatment significantly mitigated the decrease in cell viability under conditions of SD.	Cell Line:	Ovarian cancer OVCAR-3 and SKOV3 cells	Concentration:	1 μM
Cell Line:	Ovarian cancer OVCAR-3 and SKOV3 cells												
Concentration:	0, 0.2, 0.5, and 1 μM												
Incubation Time:	0, 24, 48, 72, and 96 hours												
Result:	Inhibited the decrease in cell viability induced by SD in a dose-dependent manner. 1 μM 12-HETE treatment significantly mitigated the decrease in cell viability under conditions of SD.												
Cell Line:	Ovarian cancer OVCAR-3 and SKOV3 cells												
Concentration:	1 μM												

Incubation Time:	
Result:	Led to increased levels of NF- κ B p65 phosphorylation. Caused a significant increase in the protein levels of nuclear NF- κ B p65, which was accompanied by decreased levels of NF- κ B p65 in the cytoplasm.

REFERENCES

- [1]. Qian Liu, et al. 12-HETE facilitates cell survival by activating the integrin-linked kinase/NF- κ B pathway in ovarian cancer. *Cancer Manag Res.* 2018 Nov 16;10:5825-5838.
- [2]. Benedetta Porro, et al. Analysis, physiological and clinical significance of 12-HETE: a neglected platelet-derived 12-lipoxygenase product. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2014 Aug 1;964:26-40.
- [3]. Aidan J Hampson, et al. 12-hydroxyeicosatetraenoate (12-HETE) attenuates AMPA receptor-mediated neurotoxicity: evidence for a G-protein-coupled HETE receptor. *J Neurosci.* 2002 Jan 1;22(1):257-64.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA