


Temafloxacin hydrochloride

Cat. No.:	HY-113595
CAS No.:	105784-61-0
Molecular Formula:	C ₂₁ H ₁₉ ClF ₃ N ₃ O ₃
Molecular Weight:	453.84
Target:	Antibiotic; Bacterial
Pathway:	Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Temafloxacin (TMFX) hydrochloride is an orally active quinolone broad-spectrum antibacterial agent. Temafloxacin hydrochloride is well tolerated in lower respiratory and genitourinary tract infections ^{[1][2]} .								
IC₅₀ & Target	Quinolone								
In Vitro	<p>Temafloxacin hydrochloride (0-64 µg/mL; 18-24 h) shows good antibacterial activity for gram-positive/negative bacteria, with MIC ranges of <0.004-0.5, 0.5-2 and 0.06-0.25 µg/mL for E. coli, P. aeruginosa, and S. aureus, respectively^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Viability Assay^[1]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>E. coli (16 strains), P. aeruginosa (13 strains), and S. aureus (17 strains)</td> </tr> <tr> <td>Concentration:</td> <td>0-64 µg/mL</td> </tr> <tr> <td>Incubation Time:</td> <td>18-24 h</td> </tr> <tr> <td>Result:</td> <td>Inhibited E. coli (16 strains), P. aeruginosa, and S. aureus with MIC ranges of <0.004-0.5 (MIC 90%=0.06, =0.06), 0.5-2 (MIC 90%=1, MIC 50%=1) and 0.06-0.25 µg/mL (MIC 90%=0.125, MIC 50%=0.125). MIC 90% and 50% means MIC for 90% and 50% of the isolates (unit: µg/mL).</td> </tr> </table>	Cell Line:	E. coli (16 strains), P. aeruginosa (13 strains), and S. aureus (17 strains)	Concentration:	0-64 µg/mL	Incubation Time:	18-24 h	Result:	Inhibited E. coli (16 strains), P. aeruginosa, and S. aureus with MIC ranges of <0.004-0.5 (MIC 90%=0.06, =0.06), 0.5-2 (MIC 90%=1, MIC 50%=1) and 0.06-0.25 µg/mL (MIC 90%=0.125, MIC 50%=0.125). MIC 90% and 50% means MIC for 90% and 50% of the isolates (unit: µg/mL).
Cell Line:	E. coli (16 strains), P. aeruginosa (13 strains), and S. aureus (17 strains)								
Concentration:	0-64 µg/mL								
Incubation Time:	18-24 h								
Result:	Inhibited E. coli (16 strains), P. aeruginosa, and S. aureus with MIC ranges of <0.004-0.5 (MIC 90%=0.06, =0.06), 0.5-2 (MIC 90%=1, MIC 50%=1) and 0.06-0.25 µg/mL (MIC 90%=0.125, MIC 50%=0.125). MIC 90% and 50% means MIC for 90% and 50% of the isolates (unit: µg/mL).								
In Vivo	<p>Temafloxacin hydrochloride (6.25, 25, 100 mg/kg; p.o.; single) shows good inhibitory activity to murine pyelonephritis^[1]. Temafloxacin hydrochloride (100 mg/kg; p.o. or s.c.; single) shows rapid gastrointestinal absorption, and has excellent tissue and body fluid penetration and concentration (except for central nervous system (CNS))^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>Female CF-1 mice (20-25 g) (murine pyelonephritis model)^[1].</td> </tr> <tr> <td>Dosage:</td> <td>6.25, 25, 100 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>Orally; single.</td> </tr> <tr> <td>Result:</td> <td>Reduced the number of viable bacteria in the kidneys of mice.</td> </tr> </table>	Animal Model:	Female CF-1 mice (20-25 g) (murine pyelonephritis model) ^[1] .	Dosage:	6.25, 25, 100 mg/kg	Administration:	Orally; single.	Result:	Reduced the number of viable bacteria in the kidneys of mice.
Animal Model:	Female CF-1 mice (20-25 g) (murine pyelonephritis model) ^[1] .								
Dosage:	6.25, 25, 100 mg/kg								
Administration:	Orally; single.								
Result:	Reduced the number of viable bacteria in the kidneys of mice.								

Animal Model:	Female CF-1 mice (20-25 g) ^[1] .			
Dosage:	100 mg/kg			
Administration:	Subcutaneously or orally; single.			
Result:	Pharmacokinetic Parameters of Temafloxacin hydrochloride in Female CF-1 mice ^[1] .			
	C_{max} (µg/mL)	AUC (µg/mL·h)	$T_{1/2}$ (h)	% Urinary recovery
SC (100 mg/kg)	25.2	86.6	3.4	25.3
PO (100 mg/kg)	13.5	57.4	1.3	9.1

REFERENCES

[1]. Hardy DJ, et al. Comparative antibacterial activities of temafloxacin hydrochloride (A-62254) and two reference fluoroquinolones. *Antimicrob Agents Chemother.* 1987 Nov;31(11):1768-74.

[2]. Pankey GA. Temafloxacin: an overview. *Am J Med.* 1991 Dec 30;91(6A):166S-172S.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA