


CGI-1746

Cat. No.:	HY-11999		
CAS No.:	910232-84-7		
Molecular Formula:	C ₃₄ H ₃₇ N ₅ O ₄		
Molecular Weight:	579.69		
Target:	Btk; Autophagy		
Pathway:	Protein Tyrosine Kinase/RTK; Autophagy		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 50 mg/mL (86.25 mM)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	1.7251 mL	8.6253 mL	17.2506 mL
	5 mM	0.3450 mL	1.7251 mL	3.4501 mL
	10 mM	0.1725 mL	0.8625 mL	1.7251 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2.5 mg/mL (4.31 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: 2.5 mg/mL (4.31 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: ≥ 2.5 mg/mL (4.31 mM); Suspended solution

BIOLOGICAL ACTIVITY

Description

CGI-1746 is a potent and highly selective inhibitor of the Btk with IC₅₀ of 1.9 nM.

IC₅₀ & Target

IC₅₀: 1.9 nM (Btk)

In Vitro

CGI1746 is specific for Btk, with appr 1,000-fold selectivity over Tec and Src family kinases. In an ATP-free competition binding assay, the dissociation constant for Btk is 1.5 nM. CGI1746 inhibits Btk activity in a new binding mode that stabilizes

an inactive nonphosphorylated enzyme conformation. CGI1746 inhibits both auto- and transphosphorylation steps necessary for enzyme activation. CGI1746 completely inhibits anti-IgM-induced murine and human B cell proliferation, with IC_{50} s of 134 nM and 42 nM, respectively, but has no effect on anti-CD3- and anti-CD28-induced T cell proliferation. CGI1746 potently inhibits the proliferation of CD27+IgG+ B cells isolated from the tonsils of four human donors with an average IC_{50} of 112 nM. In macrophages, CGI1746 abolishes Fc γ RIII-induced TNF α , IL-1 β and IL-6 production. CGI1746 potently inhibits TNF α , IL-1 β and, to a lesser extent, IL-6 (three- to eight-fold higher IC_{50}) production in human monocytes stimulated with immobilized or soluble immune complexes^[1]. CGI-1746 does not kill cells as well as the irreversible BTK inhibitors at the same drug concentration. CGI-1746 significantly reduces phosphorylation of both the BTK-A and BTK-C proteins, indicating the auto-phosphorylation of the BTK-C isoform is inhibited in a manner similar to BTK-A. CGI-1746 does not kill LNCaP or DU145 prostate cancer cells at the same concentrations as Ibrutinib or AVL-292, but it demonstrates similar inhibition of BTK phosphorylation at tyrosine 233 in the SH3 domain^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

CGI1746 abrogates B cell-dependent arthritis. CGI1746 treatment (100 mg/kg, s.c, twice-daily dosing) results in significant inhibition (97%) of overall clinical arthritis scores. CGI1746 treatment substantially reduces TNF α , IL-1 β and IL-6, as well as MCP1 and MIP-1 α on both the mRNA and protein level in the passive anti-collagen II antibody-induced arthritis (CAIA) model. CGI1746 shows comparable efficacy to TNF α blockade and significantly reduces clinical scores, as well as joint inflammation, in mice or rats with established arthritis^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

5×10^3 DU145 cells or 10^4 LNCaP cells per well, grown on 96 well plates for 24h, are treated with 1 to 30 μ M BTK inhibitors. Cells are fixed after 72h with 2.5% formaldehyde, and stained with Hoechst 33342. Control cells are treated with DMSO. Cell images are acquired using an IN Cell Analyzer 2200 high content imaging system, with a 20X objective. At least 9 fields are imaged per single well of each experiment. Cell numbers are determined and statistics performed using IN Cell Investigator 3.4 high content image analysis software. Each experiment is replicated 3 times, and data are presented as mean \pm SD. Results are considered significant if $p < 0.05$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nat Chem Biol. 2024 Jan 11.
- Leukemia. 2021 Feb 1.
- Mol Pharmacol. 2017 Mar;91(3):208-219.
- Patent. US20190040013A1.
- J Biomol Screen. 2015 Aug;20(7):876-86.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Di Paolo, Julie A. et al. Specific Btk inhibition suppresses B cell- and myeloid cell-mediated arthritis. Nature Chemical Biology (2011), 7(1), 41-50

[2]. Kokabee L, et al. Bruton's tyrosine kinase is a potential therapeutic target in prostate cancer. Cancer Biol Ther. 2015;16(11):1604-15

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA