


## TH1834

<b>Cat. No.:</b>	HY-123604		
<b>CAS No.:</b>	2108830-08-4		
<b>Molecular Formula:</b>	C <sub>33</sub> H <sub>40</sub> N <sub>6</sub> O <sub>3</sub>		
<b>Molecular Weight:</b>	568.71		
<b>Target:</b>	Histone Acetyltransferase; Apoptosis		
<b>Pathway:</b>	Epigenetics; Apoptosis		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 10 mg/mL (17.58 mM; ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.7584 mL	8.7918 mL	17.5837 mL
	5 mM		0.3517 mL	1.7584 mL	3.5167 mL
	10 mM		0.1758 mL	0.8792 mL	1.7584 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

TH1834 is a specific Tip60 (KAT5) histone acetyltransferase (HAT) inhibitor. TH1834 induces apoptosis and increases DNA damage in breast cancer. TH1834 does not affect the activity of related histone acetyltransferase MOF. Anticancer activity<sup>[1]</sup>.

#### In Vitro

TH1834 (0-500 μM; 1 hour; MCF7 cells) treatment significantly reduces the viability of MCF7 cells<sup>[1]</sup>.  
 TH1834 (0-500 μM; 1 hour; MCF7 cells) treatment significantly increases cytotoxicity in MCF7 cells<sup>[1]</sup>.  
 TH1834 (500 μM; 1 hour; MCF7 cells) treatment induces caspase 3 activation in MCF7 cells<sup>[1]</sup>.  
 TH1834 significantly inhibits Tip60 activity in vitro and treating cells with TH1834 results in apoptosis and increased

unrepaired DNA damage in breast cancer<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Viability Assay<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	0 $\mu$ M, 0.5 $\mu$ M, 5 $\mu$ M, 50 $\mu$ M and 500 $\mu$ M
Incubation Time:	1 hour
Result:	Significantly reduced the viability of MCF7 cells.

#### Cell Cytotoxicity Assay<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	0 $\mu$ M, 0.5 $\mu$ M, 5 $\mu$ M, 50 $\mu$ M and 500 $\mu$ M
Incubation Time:	1 hour
Result:	Highly significant increase in cytotoxicity at all concentrations used.

#### Western Blot Analysis<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	500 $\mu$ M
Incubation Time:	1 hour
Result:	Marked caspase 3 activation was observed in MCF7 cells in an independent assay.

## REFERENCES

[1]. Gao C, et al. Rational design and validation of a Tip60 histone acetyltransferase inhibitor. Sci Rep. 2014 Jun 20;4:5372.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA