


## JNJ-47117096 hydrochloride

<b>Cat. No.:</b>	HY-12420		
<b>CAS No.:</b>	1610536-69-0		
<b>Molecular Formula:</b>	C <sub>21</sub> H <sub>23</sub> ClN <sub>4</sub> O <sub>2</sub>		
<b>Molecular Weight:</b>	398.89		
<b>Target:</b>	MELK; FLT3		
<b>Pathway:</b>	PI3K/Akt/mTOR; Protein Tyrosine Kinase/RTK		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 250 mg/mL (626.74 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	2.5070 mL	12.5348 mL	25.0696 mL
	5 mM	0.5014 mL	2.5070 mL	5.0139 mL
	10 mM	0.2507 mL	1.2535 mL	2.5070 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

JNJ-47117096 hydrochloride is potent and selective MELK inhibitor, with an IC<sub>50</sub> of 23 nM, also effectively inhibits Flt3, with an IC<sub>50</sub> of 18 nM.

#### IC<sub>50</sub> & Target

IC<sub>50</sub>: 23 nM (MELK), 18 nM (Flt3)<sup>[1]</sup>

#### In Vitro

JNJ-47117096 hydrochloride is potent and selective MELK inhibitor, with an IC<sub>50</sub> of 23 nM, also effectively inhibits Flt3, with an IC<sub>50</sub> of 18 nM, and slightly blocks CAMKIIδ, Mnk2, CAMKIIγ, and MLCK (IC<sub>50</sub>, 810 nM, 760 nM, 1000 nM, 1000 nM). JNJ-47117096 (MELK-T1) suppresses the proliferation of Flt3-driven Ba/F3 cell lines, with an IC<sub>50</sub> of 1.5 μM in the absence of IL-3, while no inhibitory activity is observed in the presence of IL-3. JNJ-47117096 does not inhibit the proliferation of Ba/F3 cell lines transfected with either FGFR1, FGFR3, or KDR, either in the presence or absence of IL-3<sup>[1]</sup>. JNJ-47117096 (MELK-T1, 10 μM) delays the progression of MCF-7 cells through S-phase. JNJ-47117096 inhibits MELK, and then exerts stalled replication forks and DNA double-strand breaks (DSBs). JNJ-47117096 activates the ATM-mediated DNA-damage response (DDR). JNJ-47117096 (3, 10 μM) results in a growth arrest and a senescent phenotype. Moreover, JNJ-47117096 induces a strong phosphorylation of p53, a prolonged up-regulation of p21 and a down-regulation of FOXM1 target genes<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Kinase Assay <sup>[1]</sup>

Inhibition of MELK kinase activity is measured using a radioactive filter binding assay. Briefly, each assay well contains 1.25 nM MELK (human, residues 1-340) 10  $\mu$ M ATP, 6.7 uCi/mL  $\gamma^{33}$ P-ATP, 3  $\mu$ M biotinylated ZIP-tide peptide (Biotin-KKLNRTLSFAEPG) in 30  $\mu$ L reaction buffer (25 mM Tris pH 7.5, 10 mM MgCl<sub>2</sub>, 1 mM DTT, 1 mM EGTA, 0.1% Triton X100). Kinase reactions are performed for 25 minutes at room temperature before stopping with 40  $\mu$ L 2% orthophosphoric acid. Unbound radioactivity is removed by filtering the reaction through a MAPH filter plate. The trapped <sup>33</sup>P labelled peptide is then washed twice with 200  $\mu$ L 0.5% orthophosphoric acid, 20  $\mu$ L Microscint-20 added per well and the amount of radioactivity determined by scintillation counting using a Topcount. To calculate compound IC<sub>50</sub>, semi-log serial dilutions are used to produce 8-point dose-response curves in duplicate. IC<sub>50</sub> values are then derived using the four parameter logistic fit method in GraphPad Prism 5.0<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Cell Assay <sup>[1]</sup>

Compounds (JNJ-47117096) dissolved in DMSO are sprayed into 384-well plates (100 nL/well). A suspension of Ba/F3-Flt3 cells is added (20,000 cell/well), followed by the addition of 10 ng/mL IL3. The cells are incubated for 24 h at 37°C and 5% CO<sub>2</sub>. Alamar Blue solution is added, and after 4 h incubation at 37°C, the fluorescent intensity is measured on a Fluorescence plate reader (540 nm excitation and 590 nm emission). The control experiment in the absence of IL3 is performed in the same way. To calculate compound IC<sub>50</sub>, semi-log serial dilutions are used to produce 8-point dose-response curves in duplicate. A best-fit curve is fitted by a minimum sum of squares method to the plot of %Control vs. compound concentration. From this an IC<sub>50</sub> value is calculated<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

[1]. Johnson CN, et al. Fragment-based discovery of type I inhibitors of maternal embryonic leucine zipper kinase. ACS Med Chem Lett. 2014 May 23;6(1):25-30.

[2]. Beke L, et al. MELK-T1, a small-molecule inhibitor of protein kinase MELK, decreases DNA-damage tolerance in proliferating cancer cells. Biosci Rep. 2015 Oct 2;35(6). pii: e00267.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA