## **Product** Data Sheet

## Uridine diphosphate glucuronic acid ammonium

Cat. No.: HY-125954A CAS No.: 43195-60-4

 $\begin{tabular}{lll} \textbf{Molecular Formula:} & $C_{15}H_{22}N_2O_{18}P_2.xNH_3$ \\ \hline \textbf{Target:} & Endogenous Metabolite \\ \hline \textbf{Pathway:} & Metabolic Enzyme/Protease \\ \hline \end{tabular}$ 

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

## **BIOLOGICAL ACTIVITY**

Uridine diphosphate glucuronic acid (UDP-GlcA) ammonium is a cofactor that is formed by the catalytic activity of UDP-glucose dehydrogenase. Uridine diphosphate glucuronic acid (ammonium) is a central precursor in sugar nucleotide biosynthesis and common substrate for C4-epimerases and decarboxylases releasing UDP-galacturonic acid (UDP-GalA) and UDP-pentose products, respectively. Uridine diphosphate glucuronic acid (ammonium), as a glucuronic acid donor, can be used for for the research of the conjugation of bilirubin in the endoplasmic recticulum<sup>[1]</sup>.

IC<sub>50</sub> & Target Human Endogenous Metabolite

## **REFERENCES**

[1]. Annika J E Borg, et al. Mechanistic characterization of UDP-glucuronic acid 4-epimerase. FEBS J. 2021 Feb;288(4):1163-1178.

[2]. DUTTON GJ. Uridine diphosphate glucuronic acid as glucuronyl donor in the synthesis of ester, aliphatic and steroid glucuronides. Biochem J. 1956 Dec;64(4):693-701.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA