


4'-Methoxychalcone

Cat. No.:	HY-128400		
CAS No.:	959-23-9		
Molecular Formula:	C ₁₆ H ₁₄ O ₂		
Molecular Weight:	238.28		
Target:	PARP		
Pathway:	Cell Cycle/DNA Damage; Epigenetics		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (209.84 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	4.1967 mL	20.9837 mL	41.9674 mL
		5 mM	0.8393 mL	4.1967 mL	8.3935 mL
10 mM		0.4197 mL	2.0984 mL	4.1967 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (10.49 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	4'-Methoxychalcone regulates adipocyte differentiation through PPAR γ activation. 4'-Methoxychalcone modulates the expression and secretion of various adipokines in adipose tissue that are involved in insulin sensitivity ^[1] .
IC₅₀ & Target	PPAR γ ^[1]
In Vitro	<p>4'-Methoxychalcone (5 μM; 8 days) markedly increases the mRNA expression of adipogenic genes during adipocyte differentiation, PPARγ, aP2, FAS, adiponectin and GluT 4^[1].</p> <p>4'-Methoxychalcone (5 μM; 8 days) reduces the upregulated mRNA expression of IL-6, PAI-1, and MCP-1 by TNF-α in preadipocyte 3T3-L1 cells^[1].</p> <p>4'-Methoxychalcone (5 μM; 0-8 days) causes an increase in PPARγ expression during differentiation, while C/EBPβ protein expression is relatively unaffected^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

RT-PCR^[1]

Cell Line:	Preadipocyte 3T3-L1 cells
Concentration:	5 μ M
Incubation Time:	0 day; 2 days; 4 days; 6 days; 8 days
Result:	Increased the mRNA expression of all adipogenic genes, except GluT1 in adipocyte.

Western Blot Analysis^[1]

Cell Line:	Preadipocyte 3T3-L1 cells
Concentration:	5 μ M
Incubation Time:	8 days
Result:	Increased PPAR γ expression in adipocyte.

REFERENCES

[1]. Han Y, et al. Regulatory effects of 4-methoxychalcone on adipocyte differentiation through PPAR γ activation and reverse effect on TNF- α in 3T3-L1 cells. Food Chem Toxicol. 2017 Aug;106(Pt A):17-24.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA