

FITC-Dextran (MW 70000)

Cat. No.:	HY-128868E
CAS No.:	60842-46-8
Target:	Biochemical Assay Reagents
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

FITC-Dextran (MW 70000)

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 100 mg/mL (Need ultrasonic)
-----------------	--

BIOLOGICAL ACTIVITY

Description FITC-Dextran (MW 70000) is a compound belonging to the class of fluorescent dyes. It is commonly used in biomedical research as a tracer molecule to label and track cells or other biological matter. FITC-Dextran consists of fluorescein isothiocyanate (FITC) and dextran, a complex carbohydrate derived from starch. The combination of the two creates a stable fluorescent tracer that can be viewed under a microscope or quantified using specialized detection instruments.

In Vitro FITC-Dextran (MW 70000) is a fluorescent probe for fluorescein isothiocyanate (FITC) dextran (Ex=495 nm; Em=525 nm). FITC-Dextran (MW 70000) can be used as a marker to reveal heat shock-induced cell damage and to study the early and late stages of apoptosis. FITC-Dextran (MW 70000) can also be used for cell permeability studies, such as blood-brain barrier permeability and determination of the extent of blood-brain barrier disruption^{[1][2][3]}. Storage: protect from light. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs).
For intestinal barrier function assay^[5]

1. Fast mice for 4 h.
2. Orally gavage mice with FITC-Dextran MW 70000 (0.6 mg/g).
3. Measure fluorescence intensity of plasma in 4 h (excitation nm/emission 520 nm).

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Moumaris M, et al. Fluorescein isothiocyanate-dextran can track apoptosis and necrosis induced by heat shock of peripheral blood mononuclear cells and HeLa cells[J]. Open Biological Sciences Journal, 2015, 1(1).

[2]. Natarajan R, et al. Fluorescein Isothiocyanate (FITC)-Dextran Extravasation as a Measure of Blood-Brain Barrier Permeability. Curr Protoc Neurosci. 2017 Apr 10;79:9.58.1-9.58.15.

-
- [3]. Eriksson I, et al. Analysis of Lysosomal pH by Flow Cytometry Using FITC-Dextran Loaded Cells. *Methods Mol Biol.* 2017;1594:179-189.
- [4]. Okabayashi K, et al. Cdc42 activates paracellular transport in polarised submandibular gland cells. *Arch Oral Biol.* 2021 Dec;132:105276.
- [5]. Yu W, et al. ACE2 contributes to the maintenance of mouse epithelial barrier function. *Biochem Biophys Res Commun.* 2020 Dec 17;533(4):1276-1282.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA