


EST73502 hydrochloride

Cat. No.:	HY-134189A		
CAS No.:	2535970-65-9		
Molecular Formula:	C ₁₉ H ₂₇ ClF ₂ N ₂ O ₂		
Molecular Weight:	388.88		
Target:	Sigma Receptor; Opioid Receptor		
Pathway:	GPCR/G Protein; Neuronal Signaling		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (128.57 mM; Need ultrasonic)					
		Solvent Concentration	Mass	1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM		2.5715 mL	12.8574 mL	25.7149 mL
		5 mM		0.5143 mL	2.5715 mL	5.1430 mL
10 mM			0.2571 mL	1.2857 mL	2.5715 mL	
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.43 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.43 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.43 mM); Clear solution 					

BIOLOGICAL ACTIVITY

Description	EST73502 hydrochloride is a selective, orally active and blood-brain barrier (BBB) penetrant dual μ-opioid receptor (MOR) agonist and σ1 receptor (σ1R) antagonist, with K _i s of 64 nM and 118 nM for MOR and σ1R, respectively. EST73502 hydrochloride has antinociceptive activity ^[1] .
IC₅₀ & Target	Ki: 64 nM (MOR), 118 nM (σ1R) ^[1]
In Vivo	EST73502 hydrochloride (10-40 mg/kg; p.o.) shows a dose-response analgesic effect reaching a maximum of 64% and an EC

50 of 14 mg/kg in the paw pressure test in CD1 male mice^[1].

EST73502 hydrochloride (5 mg/kg; i.p.; twice a day; for 10 days) attenuates partial sciatic nerve ligation (PSNL)-induced mechanical allodynia in male CD1 mice^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male CD1 mice, PSNL model ^[1]
Dosage:	5 mg/kg
Administration:	Intraperitoneal injection, twice a day, for 10 days
Result:	Attenuated the expression of mechanical allodynia induced by PSNL, reaching a maximal effect of 56%.

REFERENCES

[1]. Mónica García, et al. Discovery of EST73502, a Dual μ -Opioid Receptor Agonist and σ 1 Receptor Antagonist Clinical Candidate for the Treatment of Pain. J Med Chem. 2020 Oct 16.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA