


NHC-triphosphate tetrasodium

Cat. No.:	HY-135867A
Molecular Formula:	C ₉ H ₁₂ N ₃ Na ₄ O ₁₅ P ₃
Molecular Weight:	587.08
Target:	Endogenous Metabolite; Enterovirus; HCV; Topoisomerase
Pathway:	Metabolic Enzyme/Protease; Anti-infection; Cell Cycle/DNA Damage
Storage:	-20°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (170.33 mM; Need ultrasonic)

Solvent	Mass	Concentration		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	1.7033 mL	8.5167 mL	17.0335 mL
	5 mM	0.3407 mL	1.7033 mL	3.4067 mL
	10 mM	0.1703 mL	0.8517 mL	1.7033 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

NHC-triphosphate tetrasodium is an active phosphorylated intracellular metabolite of β-d-N4-Hydroxycytidine (NHC) (HY-125033) as a triphosphate form^[1]. NHC-triphosphate tetrasodium is a weak alternative substrate for the viral polymerase and can be incorporated into HCV replicon RNA^{[1][2]}.

In Vitro

In an intracellular metabolism assay, HCV replicon cells are treated with 10 μM ³H-labeled NHC, and intracellular nucleotide levels are determined after 1, 2 and 8 hours incubations. NHC is rapidly converted into the mono-, di-, and triphosphate forms, and NHC-TP reaches up to 71.12 pM after 8 hours^[1].

NHC-triphosphate (NHC-TP) (5-40 μM) absence leads to full-length polymerization products, it can be a weak alternative substrate. In addition, incorporation of NHC-TP instead of CTP increases the molecular weight of the polymerization product by 16 (one extra oxygen) for each event and an obvious electrophoretic shift is observed in cell-free HCV NS5B polymerization reactions^[1].

Huh-7 cells are incubated with (10-50 μM; 4 h) NHC or a McGuigan phosphoramidate prodrug of NHC. Intracellular levels of the parental compounds and phosphorylated metabolites are measured using LC-MS/MS. Small amounts of NHC-monophosphate (MP) and NHC-diphosphate (DP) can be observed, while NHC-triphosphate (HY-135867) remains the most abundant metabolite^[2].

NHC-triphosphate (NHC-TP) metabolite may directly target the viral polymerase and behave as a nonobligate chain terminator. It plays a prominent role in inhibiting early negative-strand RNA synthesis, either through chain termination or

mutagenesis, which may in turn interfere with correct replicase complex formation.
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nucleic Acids Res. 2021 Jan 8;49(D1):D1113-D1121.
- ACS Sens. 2022 May 27;7(5):1564-1571.
- Commun Biol. 2022 Feb 22;5(1):154.
- J Biol Chem. 2021 May 11;100770.
- ACS Bio Med Chem Au. October 24, 2022.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Stuyver LJ, et al. Ribonucleoside analogue that blocks replication of bovine viral diarrhea and hepatitis C viruses in culture. *Antimicrob Agents Chemother.* 2003 Jan;47(1):244-54.

[2]. Maryam Ehteshami, et al. Characterization of β -d- N4-Hydroxycytidine as a Novel Inhibitor of Chikungunya Virus.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA