TRULI

Cat. No.:	HY-138489			
CAS No.:	1424635-83-5			
Molecular Formula:	C ₁₈ H ₁₄ N ₄ OS			
Molecular Weight:	334.39			
Target:	YAP			
Pathway:	Stem Cell/V	Vnt		
Storage:	Powder	-20°C	3 years	
		4°C	2 years	
	In solvent	-80°C	1 year	
		-20°C	6 months	

®

MedChemExpress

SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (149.53 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg	
		1 mM	2.9905 mL	14.9526 mL	29.9052 mL	
		5 mM	0.5981 mL	2.9905 mL	5.9810 mL	
		10 mM	0.2991 mL	1.4953 mL	2.9905 mL	
	Please refer to the so	lubility information to select the app	propriate solvent.			
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.48 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.48 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (6.22 mM); Clear solution					

BIOLOGICAL ACTIV	
Description	TRULI (Lats-IN-1) is a potent and ATP-competitive inhibitor of Lats1 and Lats2 kinases. TRULI promotes Yap-dependent proliferation in postmitotic mammalian tissues ^[1] .
In Vitro	The IC ₅₀ for TRULI increases with the ATP concentration ^[1] . TRULI (10 μM; 24 hours) interferes with the ability of Lats kinases to phosphorylate Yap, with an EC ₅₀ of 510 nM ^[1] . TRULI causes Yap-dependent proliferation of murine supporting cells in the inner ear, murine cardiomyocytes, and human Müller glia in retinal organoids ^[1] .

TRULI fosters both the O MCE has not independe	G1-S and G2-M checkpoint transitions and yields supporting cells capable of transdifferentiation ^[1] antly confirmed the accuracy of these methods. They are for reference only.
Cell Proliferation Assay	[1]
Cell Line:	MCF 10A cells
Concentration:	10 μΜ
Incubation Time:	24 hours
Result:	Decreased the phosphorylation of Yap at residue S127.

CUSTOMER VALIDATION

- EMBO J. 2023 Jan 2;e112184.
- Oncogene. 2023 Aug 17.
- PLoS Pathog. 2023 Mar 27;19(3):e1011272.
- Cells. 2023 May 29, 12(11), 1503.
- Sci Rep. 2024 Feb 18;14(1):3993.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Nathaniel Kastan, et al. Small-molecule inhibition of Lats kinases promotes Yap-dependent proliferation in postmitotic mammalian tissues. bioRxiv 2020.02.11.944157.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA