

Product Data Sheet

dGTP-15N₅,d₁₄ dilithium

Cat. No.: HY-138616S2 $\label{eq:hybrid} \mbox{Molecular Formula:} \qquad \mbox{$C_{10}D_{14}Li_2^{15}N_5O_{13}P_3$}$


Molecular Weight: 538.1

Target: Isotope-Labeled Compounds; DNA/RNA Synthesis; Nucleoside Antimetabolite/Analog D

Pathway: Others; Cell Cycle/DNA Damage

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.



BIOLOGICAL ACTIVITY

Description	dGTP- 15 N ₅ ,d ₁₄ (2'-Deoxyguanosine-5'-triphosphate- 15 N ₅ ,d ₁₄) dilithium is deuterium and 15 N labeled dGTP (HY-138616). dGTP (2'-Deoxyguanosine-5'-triphosphate), a guanosine nucleotide, can be used in deoxyribonucleic acid synthesis. Guanosine nucleotides (GDP, GTP, dGDP, and dGTP) are highly susceptible to oxidative damage to 8-oxo-GDP (8-O-GDP), 8-O-GGTP, 8-O-GTP, and 8-O-dGTP.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

 $[1]. \ Russak \ EM, et \ al. \ Impact \ of \ Deuterium \ Substitution \ on \ the \ Pharmacokinetics \ of \ Pharmaceuticals. \ Ann \ Pharmacother. \ 2019 \ Feb; 53(2): 211-216.$

[2]. Kapoor I, et al. Nucleoside Diphosphate Kinase Escalates A-to-C Mutations in MutT-Deficient Strains of Escherichia coli. J Bacteriol. 2019;202(1):e00567-19. Published 2019 Dec 6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA