


Azido-PEG1-C2-acid

Cat. No.:	HY-140009
CAS No.:	1393330-34-1
Molecular Formula:	C ₅ H ₉ N ₃ O ₃
Molecular Weight:	159.14
Target:	PROTAC Linkers
Pathway:	PROTAC
Storage:	-20°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (628.38 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	6.2838 mL	31.4189 mL	62.8377 mL
	5 mM	1.2568 mL	6.2838 mL	12.5676 mL
	10 mM	0.6284 mL	3.1419 mL	6.2838 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Azido-PEG1-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs^[1]. Azido-PEG1-C2-acid is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.

IC₅₀ & Target

PEGs

In Vitro

PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBiomedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA