Proteins

Screening Libraries

Inhibitors

Bis-aminooxy-PEG1

Cat. No.: HY-140407 CAS No.: 93460-33-4 Molecular Formula: $C_4 H_{12} N_2 O_3$ Molecular Weight: 136.15

Target: **PROTAC Linkers**

Pathway: PROTAC

Storage: -20°C, protect from light

* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

$$H_2N^O O NH_2$$

Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

H₂O: 100 mg/mL (734.48 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	7.3448 mL	36.7242 mL	73.4484 mL
	5 mM	1.4690 mL	7.3448 mL	14.6897 mL
	10 mM	0.7345 mL	3.6724 mL	7.3448 mL

Please refer to the solubility information to select the appropriate solvent.

DIO		CAL	. ACT	WITV
DIU	LUGI	CAL	. AC 11	IVIII

Description	Bis-aminooxy-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .
IC ₅₀ & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com