


Cbz-NH-PEG3-C2-acid

Cat. No.:	HY-140488	
CAS No.:	1310327-18-4	
Molecular Formula:	C ₁₇ H ₂₅ NO ₇	
Molecular Weight:	355.38	
Target:	PROTAC Linkers	
Pathway:	PROTAC	
Storage:	Pure form	-20°C 3 years
	In solvent	-80°C 6 months
		-20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (281.39 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions			1 mg	5 mg
		1 mM		2.8139 mL	14.0694 mL
		5 mM		0.5628 mL	2.8139 mL
10 mM			0.2814 mL	1.4069 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.03 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.03 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.03 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Cbz-NH-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .
IC ₅₀ & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA