RedChemExpress

Product Data Sheet

Biotin-PEG-Biotin (MW 5000)

Cat. No.:	HY-140656D	
Target:	PROTAC Linkers	
Pathway:	PROTAC	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	$\overset{H}{\underset{O}{\overset{O}{\longrightarrow}}} \overset{G}{\underset{H}{\overset{H}{\longrightarrow}}} \overset{H}{\underset{O}{\overset{O}{\longrightarrow}}} \overset{H}{\underset{O}{\overset{H}{\longrightarrow}}} \overset{H}{\underset{O}{\overset{H}{\overset{H}{\longrightarrow}}} \overset{H}{\underset{O}{\overset{H}{\overset{H}{\longrightarrow}}} \overset{H}{\underset{O}{\overset{H}{\overset{H}{\overset{H}{\overset{H}{\overset{H}{\overset{H}{\overset{H}{\overset$

BIOLOGICAL ACTIVITY		
DIOLOGICAL ACTIVITY		
Description	Biotin-PEG-Biotin (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA